

CHINAZOLINE DERIVATIVES AS AURORA KINASE INHIBITORS

The present invention relates to quinazoline derivatives for use in the treatment of disease, in particular proliferative diseases such as cancer and in the preparation of
5 medicaments for use in the treatment of proliferative diseases, and to processes for their preparation, as well as pharmaceutical compositions containing them as active ingredient.

Cancer (and other hyperproliferative diseases) are characterised by uncontrolled cellular proliferation. This loss of the normal regulation of cell proliferation often appears to occur as the result of genetic damage to cellular pathways that control progress through the
10 cell cycle.

In eukaryotes, an ordered cascade of protein phosphorylation is thought to control the cell cycle. Several families of protein kinases that play critical roles in this cascade have now been identified. The activity of many of these kinases is increased in human tumours when compared to normal tissue. This can occur by either increased levels of expression of the
15 protein (as a result of gene amplification for example), or by changes in expression of co activators or inhibitory proteins.

The first identified, and most widely studied of these cell cycle regulators have been the cyclin dependent kinases (or CDKs). Activity of specific CDKs at specific times is essential for both initiation and coordinated progress through the cell cycle. For example, the
20 CDK4 protein appears to control entry into the cell cycle (the G0-G1-S transition) by phosphorylating the retinoblastoma gene product pRb. This stimulates the release of the transcription factor E2F from pRb, which then acts to increase the transcription of genes necessary for entry into S phase. The catalytic activity of CDK4 is stimulated by binding to a partner protein, Cyclin D. One of the first demonstrations of a direct link between cancer and
25 the cell cycle was made with the observation that the Cyclin D1 gene was amplified and cyclin D protein levels increased (and hence the activity of CDK4 increased) in many human tumours (Reviewed in Sherr, 1996, Science 274: 1672-1677; Pines, 1995, Seminars in Cancer Biology 6: 63-72). Other studies (Loda et al., 1997, Nature Medicine 3(2): 231-234; Gemma et al., 1996, International Journal of Cancer 68(5): 605-11; Elledge et al. 1996, Trends in Cell
30 Biology 6; 388-392) have shown that negative regulators of CDK function are frequently down regulated or deleted in human tumours again leading to inappropriate activation of these kinases.

More recently, protein kinases that are structurally distinct from the CDK family have been identified which play critical roles in regulating the cell cycle and which also appear to be important in oncogenesis. They include the human homologues of the *Drosophila* aurora and *S.cerevisiae* Ipl1 proteins. The three human homologues of these genes Aurora-A, Aurora-B and Aurora-C (also known as aurora2, aurora1 and aurora3 respectively) encode cell cycle regulated serine-threonine protein kinases (summarised in Adams *et al.*, 2001, Trends in Cell Biology. 11(2): 49-54). These show a peak of expression and kinase activity through G2 and mitosis. Several observations implicate the involvement of human aurora proteins in cancer. This evidence is strong for Aurora-A. The Aurora-A gene maps to chromosome 20q13, a region that is frequently amplified in human tumours including both breast and colon tumours. Aurora-A may be the major target gene of this amplicon, since Aurora-A DNA is amplified and mRNA overexpressed in greater than 50% of primary human colorectal cancers. In these tumours Aurora-A protein levels appear greatly elevated compared to adjacent normal tissue. In addition, transfection of rodent fibroblasts with human Aurora-A leads to transformation, conferring the ability to grow in soft agar and form tumours in nude mice (Bischoff *et al.*, 1998, The EMBO Journal. 17(11): 3052-3065). Other work (Zhou *et al.*, 1998, Nature Genetics. 20(2): 189-93) has shown that artificial overexpression of Aurora-A leads to an increase in centrosome number and an increase in aneuploidy, a known event in the development of cancer. Further work has shown an increase in expression of Aurora-B (Adams *et al.*, 2001, Chromsoma. 110(2):65-74) and Aurora-C (Kimura *et al.*, 1999, Journal of Biological Chemistry, 274(11): 7334-40) in tumour cells when compared to normal cells.

Importantly, it has also been demonstrated that abrogation of Aurora-A expression and function by antisense oligonucleotide treatment of human tumour cell lines (WO 97/22702 and WO 99/37788) leads to cell cycle arrest and exerts an antiproliferative effect in these tumour cell lines. Additionally, small molecule inhibitors of Aurora-A and Aurora-B have been demonstrated to have an antiproliferative effect in human tumour cells (Keen *et al.* 2001, Poster #2455, American Association of Cancer research annual meeting), as has selective abrogation of Aurora-B expression alone by siRNA treatment (Ditchfield *et al.*, 2003, Journal of Cell Biology, 161(2):267-280). This indicates that inhibition of the function of Aurora-A and/or Aurora-B will have an antiproliferative effect that may be useful in the treatment of human tumours and other hyperproliferative diseases. Further, inhibition of Aurora kinases as a therapeutic approach to these diseases may have significant advantages over targeting signalling pathways upstream of the cell cycle (e.g. those activated by growth factor receptor


-3-

tyrosine kinases such as epidermal growth factor receptor (EGFR) or other receptors). Since the cell cycle is ultimately downstream of all of these diverse signalling events, cell cycle directed therapies such as inhibition of Aurora kinases would be predicted to be active across all proliferating tumour cells, whilst approaches directed at specific signalling molecules (e.g. EGFR) would be predicted to be active only in the subset of tumour cells which express those receptors. It is also believed that significant "cross talk" exists between these signalling pathways meaning that inhibition of one component may be compensated for by another.

A number of quinazoline derivatives have been proposed hitherto for use in the inhibition of Aurora kinases. For example, WO 01/21594, WO 01/21595 and WO 01/215968 describe the use of certain phenyl-quinazoline compounds as Aurora-A kinase inhibitors, which may be useful in the treatment of proliferative diseases and WO 01/21597 discloses other quinazoline derivatives as inhibitors of Aurora-A kinase. Additionally, WO 02/00649 discloses quinazoline derivative bearing a 5-membered heteroaromatic ring where the ring is, in particular, substituted thiazole or substituted thiophene. However despite the compounds of WO 02/00649 there still exists a need for further compounds having Aurora kinase inhibitory properties.

The applicants have been successful in finding a novel series of compounds which inhibit the effects of the Aurora kinases and in particular Aurora-A kinase and/or Aurora-B kinase which are thus of use in the treatment of proliferative diseases such as cancer. In particular, the compounds may be used to treat either solid or haematological tumours and more particularly colorectal, breast, lung, prostate, bladder, renal or pancreatic cancer or leukaemia or lymphoma. In addition certain aspects of the invention make them useful in the formulation of medicaments for the treatment of disease.

According to one aspect of the invention there is provided a compound of formula (I)



formula (I)

or a salt, ester or prodrug thereof;

where:

X is O or NR⁶;

R⁶ is hydrogen or C₁₋₄alkyl;

R¹ is hydrogen, halo, or -X¹R¹¹:

5 **X¹** is a direct bond, -CH₂=CH₂-, -O-, -NH-, -N(C₁₋₆alkyl)-, -C(O)-, -C(O)O-, -OC(O)-, -NHC(O)-, -N(C₁₋₆alkyl)C(O)-, -C(O)NH or -C(O)N(C₁₋₆alkyl)-;

R¹¹ is hydrogen, or a group selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkenyl, heterocyclyl, heterocyclylC₁₋₄alkyl, heterocyclylC₂₋₄alkenyl and heterocyclylC₂₋₄alkynyl which group is optionally substituted by 1 or 2 substituents

¹⁰ independently selected from halo, hydroxy, C₁₋₄alkoxy, hydroxyC₁₋₄alkyl, -NR⁹R¹⁰, -C(O)R⁹, -C(O)NR⁹R¹⁰ and -C(O)OR⁹;

R² is hydrogen, halo, nitro, cyano or -X²R¹².

X² is a direct bond, -O-, -NH-, -N(C₁₋₆alkyl)-, -OC(O)- or -C(O)O-;

R¹² is hydrogen, or a group selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₆cycloalkyl,

15 C₃₋₆cycloalkenyl, aryl, arylC₁₋₄alkyl, arylC₂₋₄alkenyl, arylC₂₋₄alkynyl, heterocyclyl, heterocyclylC₁₋₄alkyl, heterocyclylC₂₋₄alkenyl and heterocyclylC₂₋₄alkynyl, which group is optionally substituted by 1, 2 or 3 substituents independently selected from, halo, hydroxy, C₁₋₄alkyl, C₁₋₄alkoxy, -NR¹⁵R¹⁶, -NHC(O)NR¹⁵R¹⁶, -C(O)R¹⁵ and -C(O)OR¹⁵;

R³ is hydrogen, halo or -X³R¹³;

²⁰ X³ is a direct bond, -CH₂=CH₂-, -O-, -NH-, -N(C₁₋₆alkyl)-, -C(O)-, -C(O)O-, -OC(O)-, -NHC(O)-, -N(C₁₋₆alkyl)C(O)-, -C(O)NH- or -C(O)N(C₁₋₆alkyl)-;

R¹³ is hydrogen, or a group selected from C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkenyl, aryl, arylC₁₋₄alkyl, arylC₂₋₄alkenyl, arylC₂₋₄alkynyl, heterocyclyl,

heterocyclylC₁₋₄alkyl, heterocyclylC₂₋₄alkenyl and heterocyclylC₂₋₄alkynyl which group is

25 optionally substituted by 1 or 2 substituents independently selected from

$-\text{NR}^7\text{R}^8$, $-\text{C}(\text{O})\text{NR}^7\text{R}^8$, halo, hydroxy, C_{1-4} alkyl, C_{1-4} alkoxy, hydroxy C_{1-4} alkyl, hydroxy C_{1-4} alkylcarbonyl, C_{1-4} alkylcarbonyl, amino C_{1-4} alkylcarbonyl,

C₁₋₄alkylaminoC₁₋₄alkylcarbonyl and bis(C₁₋₄alkyl)aminoC₁₋₄alkylcarbonyl;

R⁷ and **R⁸** are independently selected from hydrogen, heterocyclyl, heterocyclylC₁₋₄alkyl, C₁-

30 4alkylheterocyclylC₁₋₄alkyl, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₄alkoxyC₁₋₆alkyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkylC₁₋₄alkyl, hydroxyC₃₋₆cycloalkyl, hydroxyC₁₋₄alkylC₃₋₆cycloalkyl, hydroxyC₁₋₄alkylC₃₋₆cycloalkylC₁₋₄alkyl, hydroxyC₃₋₆cycloalkylC₁₋₄alkyl, C₁₋₄alkoxyC₃₋₆cycloalkyl, C₁₋

-5-

4alkoxyC₃₋₆cycloalkylC₁₋₄alkyl, haloC₁₋₆alkyl, haloC₃₋₆cycloalkyl, haloC₃₋₆cycloalkylC₁₋₄alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, cyanoC₁₋₄alkyl, aminoC₁₋₆alkyl, C₁₋₄alkylaminoC₁₋₆alkyl, bis(C₁₋₄alkyl)aminoC₁₋₆alkyl, hydroxyC₁₋₄alkoxyC₁₋₄alkyl, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkylcarbonyl, aminoC₁₋₄alkylcarbonyl, C₁₋₄alkylaminoC₁₋₄alkylcarbonyl and bis(C₁₋₄alkyl)aminoC₁₋₄alkylcarbonyl;

or R⁷ and R⁸ together with the nitrogen to which they are attached form a heterocyclic ring which ring is monocyclic or bicyclic and comprises 4 to 7 ring atoms of which one is nitrogen and of which another is optionally selected from N, NH, O, S, SO and SO₂, and which ring is optionally substituted on carbon or nitrogen by 1 or 2 substituents independently selected from

C₁₋₄alkyl, hydroxy, C₁₋₄alkoxy, hydroxyC₁₋₄alkyl, C₁₋₄alkoxyC₁₋₄alkyl, hydroxyC₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkoxyC₁₋₄alkoxy, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkylcarbonyl, aminoC₁₋₄alkylcarbonyl, C₁₋₄alkylaminoC₁₋₄alkylcarbonyl and bis(C₁₋₄alkyl)aminoC₁₋₄alkylcarbonyl, and where a ring -CH₂- is optionally replaced with -C(O)-;

R⁴ is selected from hydrogen, halo or -X⁴R¹⁴;

X⁴ is a direct bond, -O-, -NH- or -N(C₁₋₆alkyl)-;

R¹⁴ is selected from hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl and C₂₋₆alkynyl;

R⁵ is aryl or heteroaryl optionally substituted by 1, 2 or 3 substituents independently selected from halo, hydroxy, cyano, nitro, amino, C₁₋₄alkylamino, bis(C₁₋₄alkyl)amino, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, -C(O)NHR¹⁷, -NHC(O)R¹⁸, -SR¹⁷, -S(O)R¹⁷ and

-S(O)OR¹⁷;

R⁹, R¹⁰, R¹⁵ and R¹⁶ are independently selected from hydrogen, C₁₋₆alkyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkylC₁₋₄alkyl, hydroxyC₁₋₆alkyl, haloC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₄alkylaminoC₁₋₆alkyl and bis(C₁₋₄alkyl)aminoC₁₋₆alkyl;

or R⁹ and R¹⁰ together with the nitrogen to which they are attached form a heterocyclic ring

which ring is monocyclic or bicyclic and comprises 4 to 7 ring atoms of which one is nitrogen and of which another is optionally selected from N, NH, O, S, SO and SO₂, and which ring is optionally substituted on carbon or nitrogen by 1 or 2 substituents independently selected from C₁₋₄alkyl, hydroxy, C₁₋₄alkoxy, hydroxyC₁₋₄alkyl, C₁₋₄alkoxyC₁₋₄alkyl, hydroxyC₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkoxyC₁₋₄alkoxy, hydroxyC₁₋₄alkylcarbonyl, C₁₋₄alkylcarbonyl, aminoC₁₋₄alkylcarbonyl, C₁₋₄alkylaminoC₁₋₄alkylcarbonyl and bis(C₁₋₄alkyl)aminoC₁₋₄alkylcarbonyl,


and where a ring -CH₂- is optionally replaced with -C(O)-;

-6-

R^{17} and R^{18} are independently selected from hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-4} alkenyl and C_{2-4} alkynyl.

As a further aspect a compound of formula (I) or a pharmaceutically acceptable salt thereof is provided.

5 In a further aspect the invention provides a compound of formula (IA)



formula (IA)

or a salt or ester thereof

where X, X^1 , X^2 , X^3 , R^4 and R^5 are as defined in relation to formula (I) and

10 $R^{1'}$ is hydrogen, halo, or $-X^1R^{11'}$;

$R^{11'}$ is hydrogen, phosphonooxy or a group selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, heterocyclyl, heterocyclyl C_{1-4} alkyl, heterocyclyl C_{2-4} alkenyl and heterocyclyl C_{2-4} alkynyl which group is optionally substituted by 1 or 2 substituents independently selected from halo, hydroxy, phosphonooxy, C_{1-4} alkoxy, hydroxy C_{1-4} alkyl,

15 phosphonooxy C_{1-4} alkyl, $-NR^9R^{10'}$, $-C(O)R^9$, $-C(O)NR^9R^{10'}$ and $-C(O)OR^9$;

$R^{2'}$ is hydrogen, halo, nitro, cyano or $-X^2R^{12'}$;

$R^{12'}$ is hydrogen, phosphonooxy or a group selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, aryl, aryl C_{1-4} alkyl, aryl C_{2-4} alkenyl, aryl C_{2-4} alkynyl, heterocyclyl, heterocyclyl C_{1-4} alkyl, heterocyclyl C_{2-4} alkenyl and heterocyclyl C_{2-4} alkynyl,

20 which group is optionally substituted by 1, 2 or 3 substituents independently selected from halo, hydroxy, phosphonooxy, C_{1-4} alkyl, C_{1-4} alkoxy, $-NR^{15'}R^{16'}$, $-NHC(O)NR^{15'}R^{16'}$, $-C(O)R^{15'}$ and $-C(O)OR^{15'}$;

$R^{3'}$ is hydrogen, halo or $-X^3R^{13'}$;

$R^{13'}$ is hydrogen, phosphonooxy or a group selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl,

25 C_{3-6} cycloalkyl, C_{3-6} cycloalkenyl, aryl, aryl C_{1-4} alkyl, aryl C_{2-4} alkenyl, aryl C_{2-4} alkynyl, heterocyclyl, heterocyclyl C_{1-4} alkyl, heterocyclyl C_{2-4} alkenyl and heterocyclyl C_{2-4} alkynyl which group is optionally substituted by 1 or 2 substituents independently selected from $-NR^{7'}R^8$,

-7-

$-C(O)NR^{7'}R^{8'}$, halo, hydroxy, phosphonooxy, C_{1-4} alkyl, C_{1-4} alkoxy, hydroxy C_{1-4} alkyl, phosphonooxy C_{1-4} alkyl, hydroxy C_{1-4} alkylcarbonyl, phosphonooxy C_{1-4} alkylcarbonyl, C_{1-4} alkylcarbonyl, amino C_{1-4} alkylcarbonyl, C_{1-4} alkylamino C_{1-4} alkylcarbonyl and bis(C_{1-4} alkyl)amino C_{1-4} alkylcarbonyl;

- 5 $R^{7'}$ and $R^{8'}$ are independently selected from hydrogen, heterocyclyl, heterocyclyl C_{1-4} alkyl, C_{1-4} alkylheterocyclyl C_{1-4} alkyl, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, phosphonooxy C_{1-6} alkyl, C_{1-4} alkoxy C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl C_{1-4} alkyl, hydroxy C_{3-6} cycloalkyl, phosphonooxy C_{3-6} cycloalkyl, hydroxy C_{1-4} alkyl C_{3-6} cycloalkyl, phosphonooxy C_{1-4} alkyl C_{3-6} cycloalkyl, hydroxy C_{3-6} cycloalkyl C_{1-4} alkyl,
- 10 phosphonooxy C_{3-6} cycloalkyl C_{1-4} alkyl, hydroxy C_{1-4} alkyl C_{3-6} cycloalkyl C_{1-4} alkyl, phosphonooxy C_{1-4} alkyl C_{3-6} cycloalkyl C_{1-4} alkyl, C_{1-4} alkoxy C_{3-6} cycloalkyl, C_{1-4} alkoxy C_{3-6} cycloalkyl C_{1-4} alkyl, halo C_{1-6} alkyl, halo C_{3-6} cycloalkyl, halo C_{3-6} cycloalkyl C_{1-4} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, cyano C_{1-4} alkyl, amino C_{1-6} alkyl, C_{1-4} alkylamino C_{1-6} alkyl, bis(C_{1-4} alkyl)amino C_{1-6} alkyl, hydroxy C_{1-4} alkoxy C_{1-4} alkyl,
- 15 phosphonooxy C_{1-4} alkoxy C_{1-4} alkyl, hydroxy C_{1-4} alkylcarbonyl, phosphonooxy C_{1-4} alkylcarbonyl, C_{1-4} alkylcarbonyl, amino C_{1-4} alkylcarbonyl, C_{1-4} alkylamino C_{1-4} alkylcarbonyl and bis(C_{1-4} alkyl)amino C_{1-4} alkylcarbonyl;
- or $R^{7'}$ and $R^{8'}$ together with the nitrogen to which they are attached form a heterocyclic ring which ring is monocyclic or bicyclic and comprises 4 to 7 ring atoms of which one is nitrogen
- 20 and of which another is optionally selected from N, NH, O, S, SO and SO_2 , and which ring is optionally substituted on carbon or nitrogen by 1 or 2 substituents independently selected from C_{1-4} alkyl, hydroxy, phosphonooxy, C_{1-4} alkoxy, hydroxy C_{1-4} alkyl, phosphonooxy C_{1-4} alkyl, C_{1-4} alkoxy C_{1-4} alkyl, hydroxy C_{1-4} alkoxy C_{1-4} alkyl, phosphonooxy C_{1-4} alkoxy C_{1-4} alkyl, C_{1-4} alkoxy C_{1-4} alkoxy, hydroxy C_{1-4} alkylcarbonyl, phosphonooxy C_{1-4} alkylcarbonyl,
- 25 C_{1-4} alkylcarbonyl, amino C_{1-4} alkylcarbonyl, C_{1-4} alkylamino C_{1-4} alkylcarbonyl and bis(C_{1-4} alkyl)amino C_{1-4} alkylcarbonyl, and where a ring $-CH_2-$ is optionally replaced with $-C(O)-$;
- $R^{9'}$, $R^{10'}$, $R^{15'}$ and $R^{16'}$ are independently selected from hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl C_{1-4} alkyl, hydroxy C_{1-6} alkyl, phosphonooxy C_{1-6} alkyl, halo C_{1-6} alkyl,
- 30 amino C_{1-6} alkyl, C_{1-4} alkylamino C_{1-6} alkyl and bis(C_{1-4} alkyl)amino C_{1-6} alkyl;
- or $R^{9'}$ and $R^{10'}$ together with the nitrogen to which they are attached form a heterocyclic ring which ring is monocyclic or bicyclic and comprises 4 to 7 ring atoms of which one is nitrogen

-8-

and of which another is optionally selected from N, NH, O, S, SO and SO₂, and which ring is optionally substituted on carbon or nitrogen by 1 or 2 substituents independently selected from C₁₋₄alkyl, hydroxy, phosphonooxy, C₁₋₄alkoxy, hydroxyc₁₋₄alkyl, phosphonooxyc₁₋₄alkyl, C₁₋₄alkoxyc₁₋₄alkyl, hydroxyc₁₋₄alkoxyc₁₋₄alkyl, phosphonooxyc₁₋₄alkoxyc₁₋₄alkyl, 5 C₁₋₄alkoxyc₁₋₄alkoxy, hydroxyc₁₋₄alkylcarbonyl, phosphonooxyc₁₋₄alkylcarbonyl, C₁₋₄alkylcarbonyl, aminoc₁₋₄alkylcarbonyl, C₁₋₄alkylaminoc₁₋₄alkylcarbonyl and bis(C₁₋₄alkyl)aminoc₁₋₄alkylcarbonyl, and where a ring -CH₂- is optionally replaced with -C(O)-;


provided that a compound of formula (IA) contains at least one phosphonooxy group.

10 In a preferred embodiment a compound of formula (IA) contains only one phosphonooxy group.

As a further aspect a compound of formula (IA) or a pharmaceutically acceptable salt thereof is provided.

Particular aspects of the invention provide a compound of formula (I) or a salt, ester or 15 prodrug thereof or a compound of formula (IA) or a salt, ester or prodrug thereof as described below.

A compound of formula (I) comprises



formula (I)

20 or a salt, ester or prodrug thereof;

where:

X is O or NR⁶;

R⁶ is hydrogen or C₁₋₄alkyl;

R¹ is hydrogen, halo, or -X¹R¹¹;

25 X¹ is a direct bond, -O-, -NH-, or -N(C₁₋₆alkyl)-;


- R^{11} is hydrogen, heterocyclyl or a group selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl and C_{3-6} cycloalkenyl where the group is optionally substituted by heterocyclyl, halo, hydroxy C_{1-4} alkoxy or $-NR^9R^{10}$;
- R^2 is hydrogen, halo, nitro, cyano or $-X^2R^{12}$;
- 5 X^2 is a direct bond, -O-, -NH- or $-N(C_{1-6}alkyl)-$;
- R^{12} is hydrogen, heterocyclyl or a group selected from aryl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl and C_{3-6} cycloalkenyl where the group is optionally substituted by aryl, heterocyclyl, halo, hydroxy or $-NR^{15}R^{16}$;
- R^3 is hydrogen, halo or $-X^3R^{13}$;
- 10 X^3 is a direct bond, $-CH_2=CH_2-$, -O-, -NH- or $-N(C_{1-6}alkyl)-$;
- R^{13} is hydrogen, heterocyclyl or a group selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl and C_{3-6} cycloalkenyl where the group is optionally substituted by $-NR^7R^8$, heterocyclyl, halo, hydroxy or C_{1-4} alkoxy;
- R^7 and R^8 are independently selected from hydrogen, heterocyclyl, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-3} alkoxy C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl C_{1-3} alkyl, hydroxy C_{3-6} cycloalkyl, hydroxy C_{1-4} alkyl C_{3-6} cycloalkyl, hydroxy C_{3-6} cycloalkyl C_{1-3} alkyl, C_{1-3} alkoxy C_{3-6} cycloalkyl, C_{1-3} alkoxy C_{3-6} cycloalkyl C_{1-3} alkyl, halo C_{1-6} alkyl, halo C_{3-6} cycloalkyl, halo C_{3-6} cycloalkyl C_{1-3} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, cyano C_{1-4} alkyl, amino C_{1-6} alkyl, C_{1-3} alkylamino C_{1-6} alkyl and bis(C_{1-3} alkyl)amino C_{1-6} alkyl;
- 15 or R^7 and R^8 together with the nitrogen to which they are attached form a heterocyclic ring which ring comprises 4 to 7 ring atoms of which one is nitrogen and of which another is optionally selected from N, NH, O, S, SO and SO_2 , and which ring is optionally substituted on carbon or nitrogen by 1 or 2 groups independently selected from C_{1-4} alkyl, hydroxy, C_{1-4} alkoxy, hydroxy C_{1-4} alkyl, hydroxy C_{1-4} alkoxy C_{1-4} alkyl and C_{1-4} alkoxy C_{1-4} alkoxy, and where a
- 25 ring $-CH_2-$ is optionally replaced with $-C(O)-$;
- R^4 is selected from hydrogen, halo or $-X^4R^{14}$;
- X^4 is a direct bond, -O-, -NH- or $-N(C_{1-6}alkyl)-$;
- R^{14} is selected from hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl and C_{2-6} alkynyl;
- R^5 is aryl or heteroaryl optionally substituted by 1, 2 or 3 substituents independently selected
- 30 from halo, hydroxy, cyano, nitro, amino, C_{1-4} alkylamino, bis(C_{1-4} alkyl)amino, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, $CONHR^{17}$, $NHCOR^{18}$ and $S(O)_pR^{19}$ where p is 0, 1 or 2;

-10-

R^9 , R^{10} , R^{15} and R^{16} are independently selected from hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl C_{1-3} alkyl, hydroxy C_{1-6} alkyl, halo C_{1-6} alkyl, amino C_{1-6} alkyl, C_{1-6} alkylamino C_{1-6} alkyl and bis(C_{1-6} alkyl)amino C_{1-6} alkyl;

R^{17} , R^{18} and R^{19} are independently selected from hydrogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{2-4} alkenyl and C_{2-4} alkynyl.

A compound of formula (IA) comprises



formula (IA)

where X, R^1 , R^2 , R^4 and R^5 are as defined in relation to formula (I) and

$R^{3'}$ is hydrogen, halo or $-X^{3'}R^{13'}$;

$X^{3'}$ is a direct bond, $-\text{CH}_2=\text{CH}_2-$, $-\text{O}-$, $-\text{NH}-$ or $-\text{N}(\text{C}_{1-6}\text{alkyl})-$;

$R^{13'}$ is a group selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-6} cycloalkyl and C_{3-6} cycloalkenyl where the group is substituted by $-\text{NR}^{7'}\text{R}^{8'}$;

$R^{7'}$ and $R^{8'}$ are independently selected from hydrogen, heterocyclyl, C_{1-6} alkyl,

phosphonooxy C_{1-6} alkyl, C_{1-3} alkoxy C_{1-6} alkyl, phosphonooxy C_{1-4} alkoxy C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl C_{1-3} alkyl, phosphonooxy C_{3-6} cycloalkyl, phosphonooxy C_{1-4} alkyl C_{3-6} cycloalkyl, phosphonooxy C_{3-6} cycloalkyl C_{1-3} alkyl, C_{1-3} alkoxy C_{3-6} cycloalkyl, C_{1-3} alkoxy C_{3-6} cycloalkyl C_{1-3} alkyl, halo C_{1-6} alkyl, halo C_{3-6} cycloalkyl, halo C_{3-6} cycloalkyl C_{1-3} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, cyano C_{1-4} alkyl, amino C_{1-6} alkyl, C_{1-3} alkylamino C_{1-6} alkyl and bis(C_{1-3} alkyl)amino C_{1-6} alkyl; provided that at least one of $R^{7'}$ and $R^{8'}$ contains a phosphonooxy substituent;

or $R^{7'}$ and $R^{8'}$ together with the nitrogen to which they are attached form a heterocyclic ring which ring comprises 4 to 7 ring atoms of which one is nitrogen and of which another is optionally selected from N, NH, O, S, SO and SO_2 , and which ring is substituted on carbon or nitrogen by 1 or 2 groups independently selected from phosphonooxy, phosphonooxy C_{1-4} alkyl and phosphonooxy C_{1-4} alkoxy C_{1-4} alkyl, and where a ring $-\text{CH}_2-$ is optionally replaced with a $-\text{C}(\text{O})-$.

-11-

In this specification the term alkyl when used either alone or as a suffix or prefix or otherwise includes straight-chain and branched-chain saturated structures comprising carbon and hydrogen atoms. References to individual alkyl groups such as propyl are specific for the straight-chain version only and references to individual branched-chain alkyl groups such as *tert*-butyl are specific for the branched chain version only. An analogous convention applies to other generic terms such as alkenyl and alkynyl.

Cycloalkyl is a monocyclic alkyl group, and cycloalkenyl and cycloalkynyl are monocyclic alkenyl and alkynyl groups respectively.

The prefix C_{m-n} in C_{m-n} alkyl and other terms (where m and n are integers) indicates the range of carbon atoms that are present in the group, for example C_{1-3} alkyl includes C_1 alkyl (methyl), C_2 alkyl (ethyl) and C_3 alkyl (propyl or isopropyl).

The term C_{m-n} alkoxy comprises $-O-C_{m-n}$ alkyl groups.

The term halo includes fluoro, chloro, bromo and iodo.

Aryl groups are aromatic carbocyclic groups which may be monocyclic or bicyclic.

Unless otherwise stated heteroaryl groups are monocyclic or bicyclic aromatic rings containing 5 to 10 ring atoms of which 1, 2, 3 or 4 ring atoms are chosen from nitrogen, sulphur or oxygen where a ring nitrogen or sulphur may be oxidised.

Heterocyclyl is a saturated, unsaturated or partially saturated, monocyclic or bicyclic ring containing 4 to 7 ring atoms of which 1, 2 or 3 ring atoms are selected from nitrogen, sulphur or oxygen, which ring may be carbon or nitrogen linked, wherein a $-CH_2-$ group is optionally replaced by a $-C(O)-$ group; wherein a ring nitrogen or sulphur atom is optionally oxidised to form the N-oxide or S-oxide(s); wherein a ring $-NH$ is optionally substituted by acetyl, formyl, methyl or mesyl; and wherein a ring is optionally substituted by 1 or 2 groups selected from C_{1-4} alkyl, C_{1-4} alkoxy, hydroxy C_{1-4} alkyl, hydroxy and halo C_{1-4} alkyl. In particular the ring is unsubstituted. When heterocyclyl is used within the definition of R^3 , in one aspect of the invention it is a saturated monocyclic ring containing 4 to 7 ring atoms of which one ring atom is nitrogen and another is optionally nitrogen or oxygen and which ring is optionally substituted by C_{1-4} alkyl, hydroxy C_{1-4} alkyl and hydroxy.

Phosphonooxy is in one aspect a group of formula $-OP(O)(OH)_2$. However the term phosphonooxy also includes salts such as those formed with alkali metal ions such as sodium or potassium ions or alkaline earth metal ions, for example calcium or magnesium ions.

This specification also makes use of several composite terms to describe groups comprising more than one functionality. Such terms are to be interpreted as is understood in

the art. For example C_{m-n} cycloalkyl C_{m-n} alkyl comprises C_{m-n} alkyl substituted by C_{m-n} cycloalkyl, and heterocyclyl C_{m-n} alkyl comprises C_{m-n} alkyl substituted by heterocyclyl.

Halo C_{m-n} alkyl is a C_{m-n} alkyl group that is substituted by 1, 2 or 3 halo substituents. Similarly, other generic terms containing halo such as halo C_{m-n} cycloalkyl and halo C_{m-n} cycloalkyl C_{m-n} alkyl groups may contain 1, 2 or 3 halo substituents.

Hydroxy C_{m-n} alkyl is a C_{m-n} alkyl group that is substituted by 1, 2 or 3 hydroxy substituents. Similarly other generic terms containing hydroxy such as hydroxy C_{m-n} cycloalkyl, hydroxy C_{m-n} cycloalkyl C_{m-n} alkyl, hydroxy C_{m-n} alkyl C_{m-n} cycloalkyl, hydroxy C_{m-n} alkyl C_{m-n} cycloalkyl C_{m-n} alkyl, hydroxy C_{m-n} alkoxy C_{m-n} alkyl and hydroxy C_{m-n} alkylcarbonyl groups may contain 1, 2 or 3 hydroxy substituents.

C_{m-n} alkoxy C_{m-n} alkyl is a C_{m-n} alkyl group that is substituted by 1, 2 or 3 C_{m-n} alkoxy substituents. Similarly other generic terms containing C_{m-n} alkoxy such as C_{m-n} alkoxy C_{m-n} cycloalkyl, C_{m-n} alkoxy C_{m-n} cycloalkyl C_{m-n} alkyl and C_{m-n} alkoxy C_{m-n} alkoxy groups may contain 1, 2 or 3 C_{m-n} alkoxy substituents.

Where optional substituents are chosen from 1 or 2 or from 1, 2, or 3 groups or substituents it is to be understood that this definition includes all substituents being chosen from one of the specified groups i.e. all substituents being the same or the substituents being chosen from two or more of the specified groups i.e. the substituents not being the same.

Unless specifically stated the bonding atom of a group may be any atom of that group so for example propyl includes prop-1-yl and prop-2-yl (isopropyl).

Compounds of the present invention have been named with the aid of computer software (ACD/Name version 6.6 or ACD Name Batch version 6.0).

Suitable values for any R group or any part or substituent for such groups include:

- | | |
|-----------------------------|---|
| for C_{1-4} alkyl: | methyl, ethyl, propyl, isopropyl, butyl, isobutyl and <i>tert</i> -butyl; |
| for C_{1-6} alkyl: | C_{1-4} alkyl, pentyl, neopentyl, dimethylbutyl and hexyl; |
| for C_{2-4} alkenyl: | vinyl, allyl and but-2-enyl; |
| for C_{2-6} alkenyl: | C_{2-4} alkenyl, 3-methylbut-2-enyl and 3-methylpent-2-enyl; |
| for C_{2-4} alkynyl: | ethynyl, propargyl and prop-1-ynyl; |
| for C_{2-6} alkynyl: | C_{2-4} alkynyl, pent-4-ynyl and 2-methylpent-4-ynyl; |
| for C_{3-6} cycloalkyl: | cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; |
| for C_{3-6} cycloalkenyl: | cyclobutenyl, cyclopentenyl, cyclohexenyl and cyclohex-1,4-dienyl; |

-13-

- for C₃₋₆cycloalkylC₁₋₄alkyl: cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, 2-cyclopropylethyl and 2-cyclobutylethyl;
- for C₁₋₄alkoxy: methoxy, ethoxy, propoxy, isopropoxy, butoxy and *tert*-butoxy;
- for C₁₋₄alkoxyC₁₋₄alkyl: methoxymethyl, 2-methoxyethyl, 3-methoxypropyl and 2-ethoxyethyl;
- for C₁₋₄alkoxyC₁₋₆alkyl: C₁₋₄alkoxyC₁₋₄alkyl, 4-methoxybutyl and 2-ethoxybutyl;
- for C₁₋₄alkoxyC₃₋₆cycloalkyl: 1-methoxycyclobutyl, 2-methoxycyclopentyl and 2-ethoxycyclopentyl;
- for C₁₋₄alkoxyC₃₋₆cycloalkylC₁₋₄alkyl: 1-methoxycyclobutylmethyl and 1-methoxycyclopentylmethyl;
- for C₁₋₄alkoxyC₁₋₄alkoxy: methoxymethoxy, 2-methoxyethoxy and 2-ethoxyethoxy;
- for hydroxyC₁₋₄alkyl: hydroxymethyl, 2-hydroxyethyl and 3-hydroxypropyl, 2-hydroxypropyl, 2-hydroxy-1-methylethyl, 2,3-dihydroxypropyl, 2-hydroxy-1,1-dimethylethyl;
- for hydroxyC₁₋₆alkyl: hydroxyC₁₋₄alkyl, 3-hydroxypentyl, 3-hydroxy-2,2-dimethylpropyl, 3-hydroxy-1,1-dimethylpropyl, 1-hydroxymethyl-2-methylpropyl and 6-hydroxyhexyl;
- for hydroxyC₃₋₆cycloalkyl: 2-hydroxycyclopropyl, 2-hydroxycyclobutyl, 2-hydroxycyclopentyl, and 4-hydroxycyclohexyl;
- for hydroxyC₃₋₆cycloalkylC₁₋₄alkyl: 2-hydroxycyclopropylmethyl and 2-hydroxycyclobutylmethyl;
- for hydroxyC₁₋₄alkylC₃₋₆cycloalkyl: 1-(hydroxymethyl)cyclopentyl and 2-(hydroxymethyl)cyclohexyl;
- for hydroxyC₁₋₄alkylC₃₋₆cycloalkylC₁₋₄alkyl: 1-(hydroxymethyl)cyclopropylmethyl;
- for hydroxyC₁₋₄alkoxyC₁₋₄alkyl: 2-(2-hydroxyethoxy)ethyl;
- for C₁₋₄alkylcarbonyl: acetyl, ethylcarbonyl and isopropylcarbonyl;
- for C₁₋₄alkoxycarbonyl: methoxycarbonyl, ethoxycarbonyl and *tert*-butoxycarbonyl;
- for C₁₋₄alkoxyC₁₋₄alkylcarbonyl: methoxymethylcarbonyl and *tert*-butoxymethylcarbonyl;
- for hydroxyC₁₋₄alkylcarbonyl: glycoloyl (hydroxymethylcarbonyl);
- for haloC₁₋₆alkyl: chloromethyl, 2-chloroethyl, 3-chloropropyl, trifluoromethyl and 3,3,3-trifluoropropyl;
- for haloC₃₋₆cycloalkyl: 2-chlorocyclopropyl and 2-chlorocyclobutyl;

-14-

- for haloC₃₋₆cycloalkylC₁₋₄alkyl: 2-chlorocyclopropylmethyl and 2-chlorocyclobutylmethyl;
- for cyanoC₁₋₄alkyl: cyanomethyl and 2-cyanoethyl;
- for aminoC₁₋₄alkyl: aminomethyl, 2-aminoethyl, 2-aminopropyl and 4-aminobutyl;
- 5 for aminoC₁₋₆alkyl: aminoC₁₋₄alkyl and 5-aminopentyl;
- for C₁₋₄alkylaminoC₁₋₆alkyl: 2-(methylamino)ethyl and 3-(ethylamino)propyl;
- for bis(C₁₋₄alkyl)aminoC₁₋₆alkyl: 2-(dimethylamino)ethyl, 2-[methyl(ethyl)amino]ethyl and 2-(diethylamino)ethyl;
- for C₁₋₄alkylamino: methylamino, ethylamino, propylamino and isopropylamino;
- 10 for bis(C₁₋₄alkyl)amino: dimethylamine, methyl(ethyl)amino and diethylamino;
- for aminoC₁₋₄alkylcarbonyl: glycyl (aminomethylcarbonyl);
- for C₁₋₄alkylaminoC₁₋₄alkylcarbonyl: N-methylglycyl;
- for bis(C₁₋₄alkyl)aminoC₁₋₄alkylcarbonyl: N,N-dimethylglycyl;
- for C₁₋₄alkanoylamino: acetylamino
- 15 for aryl: phenyl and naphthyl
- for arylC₁₋₄alkyl: benzyl, 2-phenylethyl;
- for arylC₂₋₄alkenyl: 3-phenylallyl;
- for arylC₂₋₄alkynyl: 3-phenylprop-2-ynyl;
- for heteroaryl: furyl, thienyl, pyrrolyl, pyrazolyl, pyridyl, pyrazinyl,
- 20 pyridazinyl, pyrimidinyl, quinazolinyl and quinolinyl
- for heterocyclyl: azetidiny, pyrrolidinyl, imidazolidinyl, piperidinyl, piperazinyl, azepanyl, diazepanyl, pyridyl, imidazolyl, tetrahydrofuranyl, tetrahydropyranyl, furanyl, pyranyl, tetrahydrothienyl, thienyl, tetrahydro-2H-pyranyl and morpholinyl.
- 25 for heterocyclylC₁₋₄alkyl: pyrrolidin-1-ylmethyl, 2-pyrrolidin-1-ylethyl, 2-morpholinoethyl, 3-morpholinopropyl, tetrahydrofuran-2-ylmethyl, 2-(2-oxopyrrolidin-3-yl)ethyl and 3-(3-oxopiperazin-1-yl)propyl;
- for heterocyclylC₂₋₄alkenyl: 3-pyrrolidin-3-ylallyl;
- 30 for heterocyclylC₂₋₄alkynyl: 3-pyrrolidin-2-ylprop-2-ynyl;
- for C₁₋₄alkylheterocyclylC₁₋₄alkyl: 5-methylisoxazol-3-ylmethyl;
- for phosphonooxyC₁₋₄alkyl: phosphonooxymethyl, 2-phosphonooxyethyl and 3-phosphonooxypropyl, 2-phosphonooxypropyl,

-15-

- for phosphonooxyC₁₋₆alkyl: 2-phosphonooxy-1-methylethyl, and
2-phosphonooxy-1,1-dimethylethyl;
phosphonooxyC₁₋₄alkyl and
3-phosphonooxy-1,1-dimethylpropyl, 3-
5 phosphonooxypentyl, 3-phosphonooxy-2,2-
dimethylpropyl, 1-phosphonooxymethyl-2-methylpropyl
and 6-phosphonooxyhexyl;
for phosphonooxyC₃₋₆cycloalkyl: 2-phosphonooxycyclopropyl, 2-phosphonooxycyclobutyl,
2-phosphonooxycyclopentyl and
10 4-phosphonooxycyclohexyl;
for phosphonooxyC₃₋₆cycloalkylC₁₋₄alkyl: 2-phosphonooxycyclopropylmethyl and 2-
phosphonooxycyclobutylmethyl;
for phosphonooxyC₁₋₄alkylC₃₋₆cycloalkyl: 1-(phosphonooxymethyl)cyclopentyl and 2-
(phosphonooxymethyl)cyclohexyl;
15 for phosphonooxyC₁₋₄alkylC₃₋₆cycloalkylC₁₋₄alkyl:
1-(phosphonooxymethyl)cyclopentylmethyl and
2-(phosphonooxymethyl)cyclohexylmethyl;
for phosphonooxyC₁₋₄alkoxyC₁₋₄alkyl: 2-(2-hydroxyethoxy)ethyl;
for phosphonooxyC₁₋₄alkylcarbonyl: phosphonooxymethylcarbonyl.

20 Within the present invention, it is to be understood that, insofar as certain compounds of formula (I) or formula (IA) herein defined may exist in optically active or racemic forms by virtue of one or more asymmetric carbon or sulphur atoms, the invention includes in its definition any such optically active or racemic form which possesses Aurora kinase inhibitory activity and in particular Aurora-A and/or Aurora-B kinase inhibitory activity. The synthesis
25 of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. Similarly, the above-mentioned activity may be evaluated using the standard laboratory techniques referred to herein.

30 Within the present invention it is to be understood that a compound of formula (I) or formula (IA) may exhibit the phenomenon of tautomerism and that the formulae drawings within this specification can represent only one of the possible tautomeric forms. It is to be understood that the invention encompasses any tautomeric form which has Aurora kinase

inhibitory activity and in particular Aurora-A and/or Aurora-B kinase inhibitory activity and is not to be limited merely to any one tautomeric form utilized within the formulae drawings.

It is also to be understood that certain compounds of formula (I) or formula (IA) and salts thereof can exist in solvated as well as unsolvated forms such as, for example, hydrated
5 forms. It is to be understood that the invention encompasses all such solvated forms which have Aurora kinase inhibitory activity and in particular Aurora-A and/or Aurora-B kinase inhibitory activity.

The present invention relates to the compounds of formula (I) or formula (IA) as herein defined as well as to the salts thereof. Salts for use in pharmaceutical compositions
10 will be pharmaceutically acceptable salts, but other salts may be useful in the production of the compounds of formula (I) or formula (IA) and their pharmaceutically acceptable salts. Pharmaceutically acceptable salts of the invention may, for example, include acid addition salts of compounds of formula (I) or formula (IA) as herein defined which are sufficiently basic to form such salts. Such acid addition salts include but are not limited to fumarate,
15 methanesulphonate, hydrochloride, hydrobromide, citrate and maleate salts and salts formed with phosphoric and sulphuric acid. In addition where compounds of formula (I) or formula (IA) are sufficiently acidic, salts are base salts and examples include but are not limited to, an alkali metal salt for example sodium or potassium, an alkaline earth metal salt for example calcium or magnesium, or organic amine salt for example triethylamine, ethanolamine,
20 diethanolamine, triethanolamine, morpholine, *N*-methylpiperidine, *N*-ethylpiperidine, dibenzylamine or amino acids such as lysine.

The compounds of formula (I) or formula (IA) may also be provided as *in vivo* hydrolysable esters. An *in vivo* hydrolysable ester of a compound of formula (I) or formula (IA) containing carboxy or hydroxy group is, for example a pharmaceutically acceptable ester
25 which is cleaved in the human or animal body to produce the parent acid or alcohol. Such esters can be identified by administering, for example, intravenously to a test animal, the compound under test and subsequently examining the test animal's body fluid.

Suitable pharmaceutically acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl; C₁₋₆alkanoyloxymethyl esters for example
30 pivaloyloxymethyl; phthalidyl esters; C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example

-17-

1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

Suitable pharmaceutically-acceptable esters for hydroxy include inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxymethoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include C₁₋₁₀alkanoyl, for example formyl, acetyl; benzoyl; phenylacetyl; substituted benzoyl and phenylacetyl; C₁₋₁₀alkoxycarbonyl (to give alkyl carbonate esters), for example ethoxycarbonyl; di-C₁₋₄alkylcarbamoyl and *N*-(di-C₁₋₄alkylaminoethyl)-*N*-C₁₋₄alkylcarbamoyl (to give carbamates); di-C₁₋₄alkylaminoacetyl and carboxyacetyl. Examples of ring substituents on phenylacetyl and benzoyl include aminomethyl, C₁₋₄alkylaminomethyl and di-(C₁₋₄alkyl)aminomethyl, and morpholino or piperazino linked from a ring nitrogen atom via a methylene linking group to the 3- or 4- position of the benzoyl ring. Other interesting *in vivo* hydrolysable esters include, for example, R^AC(O)OC₁₋₆alkyl-CO-, wherein R^A is for example, benzyloxy-C₁₋₄alkyl, or phenyl. Suitable substituents on a phenyl group in such esters include, for example, 4-C₁₋₄piperazino-C₁₋₄alkyl, piperazino-C₁₋₄alkyl and morpholino-C₁₋₄alkyl.

The compounds of the formula (I) may be also be administered in the form of a prodrug which is broken down in the human or animal body to give a compound of the formula (I). Examples of prodrugs include *in vivo* hydrolysable esters of a compound of the formula (I). Various forms of prodrugs are known in the art. For examples of such prodrug derivatives, see:

- a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, et al. (Academic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard p. 113-191 (1991);
- c) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);
- d) H. Bundgaard, et al., Journal of Pharmaceutical Sciences, 77, 285 (1988); and
- e) N. Kakeya, et al., Chem Pharm Bull, 32, 692 (1984).

-18-

Particular values of X, R¹, R^{1'}, R², R^{2'}, R³, R^{3'}, R⁴ and R⁵ for compounds of formula (I) and formula (IA) are as follows. Such values may be used where appropriate with any of the definitions, claims or embodiments defined herein.

In one aspect of the invention X is NR⁶. In another aspect X is NH.

5 In one aspect of the invention R⁶ is hydrogen or methyl. In another aspect R⁶ is hydrogen.

In one aspect of the invention R¹ is hydrogen or -OR¹¹. In another aspect R¹ is hydrogen.

10 In one aspect of the invention X¹ is a direct bond or -O-. In another aspect X¹ is a direct bond.

In one aspect of the invention R¹¹ is hydrogen, heterocyclyl selected from piperidinyl or pyrrolidinyl or C₁₋₄alkyl which C₁₋₄alkyl is optionally substituted by hydroxy, C₁₋₄alkoxy, amino, C₁₋₄alkylamino or bis(C₁₋₄alkyl)amino. In another aspect R¹¹ is hydrogen, C₁₋₄alkyl or C₁₋₄alkoxy. In another aspect R¹¹ is hydrogen.

15 In one aspect of the invention R² is hydrogen or -OR¹². In another aspect R² is hydrogen or methoxy. In a further aspect R² is hydrogen. In yet a further aspect R² is methoxy.

In one aspect of the invention X² is a direct bond or -O-. In another aspect X² is a direct bond. In a further aspect X² is -O-.

20 In one aspect of the invention R¹² is hydrogen, C₁₋₄alkyl, heterocyclyl or heterocyclylC₁₋₄alkyl. In another aspect R¹² is hydrogen or C₁₋₄alkyl. In another aspect of the invention R¹² is hydrogen. In a further aspect of the invention R¹² is methyl.

In one aspect of the invention R³ is -X³R¹³. In a further aspect R³ is selected from 3-chloropropoxy, 3-[2-(hydroxymethyl)pyrrolidin-1-yl]propoxy, 3-[(2-hydroxyethyl)(isobutyl)amino]propoxy, 3-[(2-hydroxyethyl)(propyl)amino]propoxy, 3-piperidin-1-ylpropoxy, 3-pyrrolidin-1-ylpropoxy, 3-(diethylamino)propoxy, 3-piperazin-1-ylpropoxy, 3-[(2-hydroxyethyl)(methyl)amino]propoxy, 3-(cyclopropylamino)propoxy, 3-[[2-(dimethylamino)ethyl](methyl)amino]propoxy, 3-(4-methylpiperazin-1-yl)propoxy, 3-(4-hydroxypiperidin-1-yl)propoxy, 3-[bis(2-hydroxyethyl)amino]propoxy, 3-ethyl(methyl)amino]propoxy, 3-[ethyl(2-hydroxyethyl)amino]propoxy, 3-[[2-(dimethylamino)ethyl](ethyl)amino]propoxy, 3-[2-(2-hydroxyethyl)piperidin-1-yl]propoxy, 3-[4-(2-hydroxyethyl)piperazin-1-yl]propoxy, 3-[(cyclopropylmethyl)amino]propoxy, 3-[4-(2-hydroxyethyl)piperidin-1-yl]propoxy, 3-[methyl(propargyl)amino]propoxy, 3-

25

30

-19-

[allyl(methyl)amino]propoxy, 3-[isobutyl(methyl)amino]propoxy, 3-(3-hydroxypiperidin-1-yl)propoxy, 3-[4-(hydroxymethyl)piperidin-1-yl]propoxy, 3-[methyl(propyl)amino]propoxy, 3-[cyclopropylmethyl(propyl)amino]propoxy, 3-{[2-(diethylamino)ethyl](methyl)amino}propoxy, 3-{[2-(diethylamino)ethyl](ethyl)amino}propoxy, 3-(4-methyl-1,4-diazepan-1-yl)propoxy, 3-[(2-hydroxyethyl)(isopropyl)amino]propoxy, 3-[cyclopropyl(2-hydroxyethyl)amino]propoxy, 3-[(2-hydroxyethyl)(2-methoxyethyl)amino]propoxy, 3-[cyclobutyl(2-hydroxyethyl)amino]propoxy, 3-[cyclopropylmethyl(2-hydroxyethyl)amino]propoxy, 3-[cyclobutylmethyl(2-hydroxyethyl)amino]propoxy, 3-[(2-hydroxy)propargylamino]propoxy, 3-[allyl(2-hydroxyethyl)amino]propoxy, 3-[(2-hydroxyethyl)neopentylamino]propoxy, 3-[(2-hydroxyethyl)(3,3,3-trifluoropropyl)amino]propoxy, 3-azetidin-3-ylpropoxy, 3-[cyclopentyl(2-hydroxyethyl)amino]propoxy, 3-[(3-hydroxy-1,1-dimethylpropyl)amino]propoxy, 3-[(2-cyanoethyl)(2-hydroxyethyl)amino]propoxy, 3-(dimethylamino)propoxy, 3-[(2-hydroxy-1,1-dimethylethyl)amino]propoxy and 3-morpholin-4-ylpropoxy. In another aspect R³ is selected from 3-[2-(hydroxymethyl)pyrrolidin-1-yl]propoxy, 3-[(2-hydroxyethyl)(isobutyl)amino]propoxy, 3-[(2-hydroxyethyl)(propyl)amino]propoxy, 3-[ethyl(2-hydroxyethyl)amino]propoxy, 3-[4-(2-hydroxyethyl)piperazin-1-yl]propoxy, 3-[4-(2-hydroxyethyl)piperidin-1-yl]propoxy, 3-[(2-hydroxyethyl)(2-methoxyethyl)amino]propoxy, 3-[cyclobutyl(2-hydroxyethyl)amino]propoxy, 3-[cyclopropylmethyl(2-hydroxyethyl)amino]propoxy and 3-[(3-hydroxy-1,1-dimethylpropyl)amino]propoxy. In yet another aspect R³ is 3-[(2-hydroxyethyl)(propyl)amino]propoxy, 3-[2-(hydroxymethyl)pyrrolidin-1-yl]propoxy, 3-morpholin-4-ylpropoxy, 3-piperidin-1-ylpropoxy, 3-pyrrolidin-1-ylpropoxy, 3-[(2-hydroxy-1,1-dimethylethyl)amino]propoxy, 3-(cyclopropylamino)propoxy, 3-[[2-(dimethylamino)ethyl](methyl)amino]propoxy, 3-[[2-(dimethylamino)ethyl](ethyl)amino]propoxy, 3-(4-methylpiperazin-1-yl)propoxy, 3-(4-hydroxypiperidin-1-yl)propoxy, 3-[ethyl(2-hydroxyethyl)amino]propoxy, 3-[4-(2-hydroxyethyl)piperazin-1-yl]propoxy, 3-piperazin-1-ylpropoxy, 3-[4-(2-hydroxyethyl)piperidin-1-yl]propoxy, 3-[4-(hydroxymethyl)piperidin-1-yl]propoxy, 3-[(2-hydroxyethyl)(isopropyl)amino]propoxy and 3-[cyclopropyl(2-hydroxyethyl)amino]propoxy. In another aspect R³ is 3-chloropropoxy. In a further aspect R³ is 3-chloropropoxy, 3-[2-(hydroxymethyl)pyrrolidin-1-yl]propoxy and 3-[(2-hydroxyethyl)(propyl)amino]propoxy.

In one aspect of the invention X³ is -CH₂=CH₂-, -O- or -NH-. In another aspect X³ is -O-.

-20-

In one aspect of the invention R^{13} is C_{1-6} alkyl substituted by $-NR^7R^8$, heterocyclyl or halo. In a further aspect of the invention R^{13} is ethyl or propyl, which ethyl or propyl are substituted by $-NR^7R^8$, heterocyclyl or halo. In yet a further aspect of the invention R^{13} is propyl substituted by chloro, $-NR^7R^8$ or a heterocyclyl selected from pyrrolidinyl, piperidinyl, 5 piperazinyl, morpholinyl, diazepanyl and azetidiny where the heterocyclyl is optionally substituted by hydroxy, methyl, hydroxymethyl or 2-hydroxyethyl. In another aspect R^{13} is propyl substituted by chloro or $-NR^7R^8$. In a further aspect R^{13} is propyl substituted by $-NR^7R^8$.

In one aspect of the invention R^7 and R^8 are independently selected from hydrogen, 10 heterocyclyl, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, hydroxy C_{1-4} alkyl C_{3-6} cycloalkyl, C_{1-4} alkoxy C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl C_{1-4} alkyl, halo C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, cyano C_{1-4} alkyl and bis(C_{1-4} alkyl)amino C_{1-6} alkyl; or R^7 and R^8 together with the nitrogen to which they are attached form a heterocyclic ring which ring comprises 4 to 7 ring atoms of which one is nitrogen and of which another is optionally NH or O and which ring is optionally substituted 15 on carbon or nitrogen by a group selected from C_{1-4} alkyl, hydroxy, hydroxy C_{1-4} alkyl and hydroxy C_{1-4} alkoxy C_{1-4} alkyl, and where a ring $-CH_2-$ is optionally replaced with $-C(O)-$. In a further aspect R^7 and R^8 are independently selected from hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *tert*-butyl, pentyl, neopentyl, hydroxymethyl, 2-hydroxyethyl, 2-hydroxy-1,1-dimethylethyl, 3-hydroxy-1,1-dimethylpropyl, methoxymethyl, 2-methoxyethyl, 20 2-ethoxyethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 3,3,3-trifluoropropyl, allyl, propargyl, 2-(dimethylamino)ethyl and 2-(diethylamino)ethyl; or R^7 and R^8 together with the nitrogen to which they are attached form a heterocyclic ring selected from pyrrolidine, piperidine, piperazine, morpholine, diazepane and azetidine which ring is optionally substituted by 25 hydroxy, methyl, hydroxymethyl or 2-hydroxyethyl. In yet another aspect R^7 and R^8 are independently selected from hydrogen, methyl, ethyl, propyl, isopropyl, isobutyl, 2-hydroxyethyl, 2-hydroxy-1,1-dimethylethyl, 3-hydroxy-1,1-dimethyl, 2-methoxyethyl, cyclopropyl, cyclobutyl, cyclopropylmethyl and 2-(dimethylamino)ethyl; or R^7 and R^8 together with the nitrogen to which they are attached form a heterocyclic ring selected from 30 pyrrolidine, piperidine, piperazine and morpholine, which the ring is optionally substituted by hydroxy, methyl, hydroxymethyl or 2-hydroxyethyl. In yet another aspect R^7 and R^8 are independently selected from hydrogen, methyl, ethyl, propyl, isopropyl, cyclopropyl, 2-hydroxyethyl, 2-hydroxy-1,1-dimethylethyl and 2-(dimethylamino)ethyl; or R^7 and R^8

-21-

together with the nitrogen to which they are attached form a heterocyclic ring selected from pyrrolidine, piperidine, piperazine and morpholine, which the ring is optionally substituted by hydroxy, methyl, hydroxymethyl or 2-hydroxyethyl. In a further aspect R^7 and R^8 are independently propyl or 2-hydroxyethyl; or R^7 and R^8 together with the nitrogen to which they
 5 are attached form pyrrolidine substituted by hydroxymethyl.

In one aspect of the invention R^4 is hydrogen.

In one aspect of the invention R^5 is aryl optionally substituted by 1 or 2 halo. In another aspect R^5 is phenyl optionally substituted by 1 or 2 fluoro or chloro. In a further aspect R^5 is phenyl optionally substituted by 1 or 2 fluoro. In yet another aspect R^5 is 2,3-
 10 difluorophenyl or 3-fluorophenyl. In another aspect R^5 is 3-fluorophenyl.

In one aspect of the invention $R^{1'}$ is hydrogen or $-OR^{11'}$. In another aspect $R^{1'}$ is hydrogen.

In one aspect of the invention $R^{11'}$ is hydrogen, heterocyclyl selected from piperidinyl or pyrrolidinyl, C_{1-4} alkyl optionally substituted by hydroxy, C_{1-4} alkoxy, amino, C_{1-4} alkylamino
 15 or bis(C_{1-4} alkyl)amino.

In one aspect of the invention $R^{2'}$ is hydrogen or $-OR^{12'}$. In another aspect $R^{2'}$ is hydrogen or methoxy.

In one aspect of the invention $R^{12'}$ is hydrogen, C_{1-4} alkyl (optionally substituted with heterocyclyl) or heterocyclyl;

20 In one aspect of the invention $R^{3'}$ is $-X^{3'}R^{13'}$. In a further aspect $R^{3'}$ is selected from 3-[propyl(2-phosphonooxyethyl)amino]propoxy, 3-(2-phosphonooxymethylpyrrolidin-1-yl)propoxy, 3-[ethyl(2-phosphonooxyethyl)amino]propoxy, 3-[(2-methoxyethyl)(2-phosphonooxyethyl)amino]propoxy, 3-[cyclobutyl(2-phosphonooxyethyl)amino]propoxy, 3-[4-(2-phosphonooxymethyl)piperazin-1-yl]propoxy and 3-[(1,1-dimethyl-3-phosphonooxypropyl)amino]propoxy. In yet another aspect $R^{3'}$ is 3-[(2-phosphonooxyethyl)(propyl)amino]propoxy, 3-[2-(phosphonooxymethyl)pyrrolidin-1-yl]propoxy, 3-morpholin-4-ylpropoxy, 3-piperidin-1-ylpropoxy, 3-pyrrolidin-1-ylpropoxy, 3-[(2-phosphonooxy-1,1-dimethylethyl)amino]propoxy, 3-(cyclopropylamino)propoxy, 3-[[2-dimethylamino)ethyl](methyl)amino]propoxy, 3-[[2-dimethylamino)ethyl](ethyl)amino]propoxy, 3-(4-methylpiperazin-1-yl)propoxy, 3-(4-phosphonooxypiperidin-1-yl)propoxy, 3-[ethyl(2-phosphonooxyethyl)amino]propoxy, 3-[4-(2-phosphonooxyethyl)piperazin-1-yl]propoxy, 3-piperazin-1-ylpropoxy, 3-[4-(2-phosphonooxyethyl)piperidin-1-yl]propoxy, 3-[4-(phosphonooxymethyl)piperidin-1-

-22-

yl]propoxy, 3-[(2-phosphonooxyethyl)(isopropyl)amino]propoxy and 3-[cyclopropyl(2-phosphonooxyethyl)amino]propoxy.

In one aspect of the invention $X^{3'}$ is $-\text{CH}_2=\text{CH}_2-$, $-\text{O}-$ or $-\text{NH}-$. In a further aspect $X^{3'}$ is $-\text{O}-$.

5 In one aspect of the invention $R^{13'}$ is C_{1-6} alkyl substituted by $-\text{NR}^{7'}\text{R}^{8'}$. In a further aspect of the invention $R^{13'}$ is propyl substituted by $-\text{NR}^{7'}\text{R}^{8'}$.

In one aspect of the invention $R^{7'}$ is selected from hydrogen, heterocyclyl, C_{1-6} alkyl, C_{1-4} alkoxy C_{1-6} alkyl, cyano C_{1-4} alkyl, C_{3-6} cycloalkyl, amino C_{1-6} alkyl, C_{1-4} alkylamino C_{1-6} alkyl and bis(C_{1-4} alkyl)amino C_{1-6} alkyl. In a further aspect $R^{7'}$ is hydrogen, methyl, ethyl, propyl,
10 isopropyl, cyclopropyl and 2-(dimethylamino)ethyl. In another aspect $R^{7'}$ is ethyl, propyl, cyclobutyl or 2-methoxyethyl.

In one aspect of the invention $R^{8'}$ is phosphonooxy C_{1-4} alkyl or phosphonooxy C_{1-4} alkyl C_{3-6} cycloalkyl. In a further aspect $R^{8'}$ is phosphonooxy C_{1-4} alkyl. In another aspect $R^{8'}$ is 2-phosphonooxyethyl or 1,1-dimethyl-2-phosphonooxyethyl.

15 In one aspect of the invention $R^{7'}$ and $R^{8'}$ together with the nitrogen to which they are attached form a heterocyclic ring selected from pyrrolidine, piperidine, piperazine and morpholine which ring is substituted on carbon or nitrogen by a group selected from phosphonooxy, phosphonooxymethyl and 2-phosphonooxyethyl.

20 A particular class of compounds is of formula (I) wherein:

X is NR^6 ;

R^6 is hydrogen or methyl;

R^1 is hydrogen or $-\text{OR}^{11}$;

R^{11} is hydrogen, heterocyclyl selected from piperidinyl or pyrrolidinyl or C_{1-4} alkyl which C_{1-4} alkyl is optionally substituted by hydroxy, C_{1-4} alkoxy, amino, C_{1-4} alkylamino or bis(C_{1-4} alkyl)amino;

25 R^2 is hydrogen or $-\text{OR}^{12}$;

R^{12} is hydrogen, C_{1-4} alkyl, heterocyclyl or heterocyclyl C_{1-4} alkyl;

R^3 is $-\text{X}^3\text{R}^{13}$;

30 X^3 is $-\text{CH}_2=\text{CH}_2-$, $-\text{O}-$ or $-\text{NH}-$;

R^{13} is C_{1-6} alkyl substituted by $-\text{NR}^{7'}\text{R}^{8'}$, heterocyclyl or halo;

R^7 and R^8 are independently selected from hydrogen, heterocyclyl, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, hydroxy C_{1-4} alkyl C_{3-6} cycloalkyl, C_{1-4} alkoxy C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{3-6} cycloalkyl C_{1-4} alkyl,

-23-

alkyl, haloC₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, cyanoC₁₋₄alkyl and bis(C₁₋₄alkyl)aminoC₁₋₆alkyl; or R⁷ and R⁸ together with the nitrogen to which they are attached form a heterocyclic ring which ring comprises 4 to 7 ring atoms of which one is nitrogen and of which another is optionally NH or O and which ring is optionally substituted on carbon or nitrogen by a group
 5 selected from C₁₋₄alkyl, hydroxy, hydroxyC₁₋₄alkyl and hydroxyC₁₋₄alkoxyC₁₋₄alkyl, and where a ring -CH₂- is optionally replaced with -C(O)-;
 R⁴ is hydrogen; and
 R⁵ is aryl optionally substituted by 1 or 2 halo;
 or a salt, ester or prodrug thereof.

10

A further class of compounds is of formula (I) wherein:

X is NH;

R¹ is hydrogen;

R² is hydrogen or methoxy;

15 R³ is -X³R¹³;

X³ is -O-;

R¹³ is propyl substituted by chloro or -NR⁷R⁸;

R⁷ and R⁸ are independently selected from hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *tert*-butyl, pentyl, neopentyl, hydroxymethyl, 2-hydroxyethyl, 2-hydroxy-1,1-
 20 dimethylethyl, 3-hydroxy-1,1-dimethylpropyl, methoxymethyl, 2-methoxyethyl, 2-ethoxyethyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 3,3,3-trifluoropropyl, allyl, propargyl, 2-(dimethylamino)ethyl and 2-(diethylamino)ethyl; or R⁷ and R⁸ together with the nitrogen to which they are attached form a heterocyclic ring selected from pyrrolidine, piperidine,
 25 piperazine, morpholine, diazepane and azetidine which ring is optionally substituted by hydroxy, methyl, hydroxymethyl or 2-hydroxyethyl;
 R⁴ is hydrogen; and
 R⁵ is 2,3-difluorophenyl or 3-fluorophenyl;
 or a salt, ester or prodrug thereof.

30

A further class of compounds is of formula (I) wherein:

X is NH;

R¹ is hydrogen;

-24-

R^2 is hydrogen or methoxy;

R^3 is $-X^3R^{13}$;

X^3 is $-O-$;

R^{13} is propyl substituted by chloro or $-NR^7R^8$;

- 5 R^7 and R^8 are independently selected from hydrogen, methyl, ethyl, propyl, isopropyl, cyclopropyl, 2-hydroxyethyl, 2-hydroxy-1,1-dimethylethyl and 2-(dimethylamino)ethyl; or R^7 and R^8 together with the nitrogen to which they are attached form a heterocyclic ring selected from pyrrolidine, piperidine, piperazine and morpholine, which the ring is optionally substituted by hydroxy, methyl, hydroxymethyl or 2-hydroxyethyl;
- 10 R^4 is hydrogen; and
- R^5 is 2,3-difluorophenyl or 3-fluorophenyl;
- or a salt, ester or prodrug thereof.

A particular class of compounds is of formula (IA) wherein:

- 15 X is NR^6 ;
- R^6 is hydrogen or methyl;
- $R^{1'}$ is hydrogen or $-OR^{11}$;
- $R^{11'}$ is hydrogen, heterocyclyl selected from piperidinyl or pyrrolidinyl, C_{1-4} alkyl optionally substituted by hydroxy, C_{1-4} alkoxy, amino, C_{1-4} alkylamino or bis(C_{1-4} alkyl)amino;
- 20 $R^{2'}$ is hydrogen or $-OR^{12}$;
- $R^{12'}$ is hydrogen, C_{1-4} alkyl (optionally substituted with heterocyclyl) or heterocyclyl;
- $R^{3'}$ is $-X^{3'}R^{13'}$;
- $X^{3'}$ is $-CH_2=CH_2-$, $-O-$ or $-NH-$;
- $R^{13'}$ is C_{1-6} alkyl substituted by $-NR^7R^8$;
- 25 $R^{7'}$ is hydrogen, methyl, ethyl, propyl, isopropyl, cyclopropyl or 2-(dimethylamino)ethyl;
- $R^{8'}$ is 2-phosphonooxyethyl or 1,1-dimethyl-2-3-phosphonooxyethyl;
- or $R^{7'}$ and $R^{8'}$ together with the nitrogen to which they are attached form a heterocyclic ring selected from pyrrolidine, piperidine, piperazine and morpholine which ring is substituted on carbon or nitrogen by a group selected from phosphonooxy, phosphonooxymethyl and 2-
- 30 phosphonooxyethyl;
- R^4 is hydrogen; and
- R^5 is aryl optionally substituted by 1 or 2 halo;
- or a salt or prodrug thereof.

Particular compounds of the invention are any one of:

- 2-(4-{[7-(3-chloropropoxy)-6-methoxyquinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)-*N*-(3-fluorophenyl)acetamide;
- 5 2-(4-{[7-(3-chloropropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)-*N*-(3-fluorophenyl)acetamide;
- (4-{[7-(3-chloropropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)-*N*-(2,3-difluorophenyl)acetamide;
- N*-(3-fluorophenyl)-2-{4-[(7-{3-[(2-hydroxyethyl)(propyl)amino]propoxy}-6-methoxyquinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- 10 *N*-(3-fluorophenyl)-2-{4-[(7-{3-[(2*S*)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy}-6-methoxyquinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- N*-(3-fluorophenyl)-2-{4-[(7-{3-[(2-hydroxyethyl)(propyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- 15 *N*-(3-fluorophenyl)-2-{4-[(7-{3-[(2*S*)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- N*-(3-fluorophenyl)-2-(4-{[7-(3-morpholin-4-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide;
- N*-(3-fluorophenyl)-2-(4-{[7-(3-piperidin-1-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide;
- 20 *N*-(3-fluorophenyl)-2-(4-{[7-(3-pyrrolidin-1-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide;
- N*-(3-fluorophenyl)-2-{4-[(7-{3-[(2-hydroxy-1,1-dimethylethyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- 25 2-[4-({7-[3-(cyclopropylamino)propoxy]quinazolin-4-yl}amino)-1*H*-1,2,3-triazol-1-yl]-*N*-(3-fluorophenyl)acetamide;
- 2-{4-[(7-{3-[(2-(dimethylamino)ethyl)(methyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}-*N*-(3-fluorophenyl)acetamide;
- N*-(3-fluorophenyl)-2-[4-({7-[3-(4-methylpiperazin-1-yl)propoxy]quinazolin-4-yl}amino)-1*H*-1,2,3-triazol-1-yl]acetamide;
- 30 1,2,3-triazol-1-yl]acetamide;
- N*-(3-fluorophenyl)-2-{4-[(7-{3-[(2*R*)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;

- N*-(3-fluorophenyl)-2-[4-({7-[3-(4-hydroxypiperidin-1-yl)propoxy]quinazolin-4-yl}amino)-1*H*-1,2,3-triazol-1-yl]acetamide;
- 2-[4-[(7-{3-[ethyl(2-hydroxyethyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl]-*N*-(3-fluorophenyl)acetamide;
- 5 *N*-(3-fluorophenyl)-2-{4-[(7-{3-[4-(2-hydroxyethyl)piperazin-1-yl]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- N*-(3-fluorophenyl)-2-(4-{[7-(3-piperazin-1-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide;
- N*-(3-fluorophenyl)-2-{4-[(7-{3-[4-(2-hydroxyethyl)piperidin-1-yl]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- 10 *N*-(3-fluorophenyl)-2-{4-[(7-{3-[4-(hydroxymethyl)piperidin-1-yl]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- N*-(3-fluorophenyl)-2-{4-[(7-{3-[(2-hydroxyethyl)(isopropyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- 15 2-[4-[(7-{3-[cyclopropyl(2-hydroxyethyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl]-*N*-(3-fluorophenyl)acetamide;
- N*-(2,3-difluorophenyl)-2-(4-{[7-(3-morpholin-4-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide;
- N*-(2,3-difluorophenyl)-2-(4-{[7-(3-piperidin-1-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide;
- 20 *N*-(2,3-difluorophenyl)-2-(4-{[7-(3-pyrrolidin-1-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide;
- N*-(2,3-difluorophenyl)-2-{4-[(7-{3-[(2-hydroxy-1,1-dimethylethyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- 25 2-[4-({7-[3-(cyclopropylamino)propoxy]quinazolin-4-yl}amino)-1*H*-1,2,3-triazol-1-yl]-*N*-(2,3-difluorophenyl)acetamide;
- N*-(2,3-difluorophenyl)-2-{4-[(7-{3-[[2-(dimethylamino)ethyl](methyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;
- 30 *N*-(2,3-difluorophenyl)-2-[4-({7-[3-(4-methylpiperazin-1-yl)propoxy]quinazolin-4-yl}amino)-1*H*-1,2,3-triazol-1-yl]acetamide;
- N*-(2,3-difluorophenyl)-2-{4-[(7-{3-[(2*R*)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;

-27-

N-(2,3-difluorophenyl)-2-[4-((7-[3-(4-hydroxypiperidin-1-yl)propoxy]quinazolin-4-yl)amino)-1*H*-1,2,3-triazol-1-yl]acetamide;

N-(2,3-difluorophenyl)-2-{4-[(7-{3-[ethyl(2-hydroxyethyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;

5 *N*-(2,3-difluorophenyl)-2-{4-[(7-{3-[4-(2-hydroxyethyl)piperazin-1-yl]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;

N-(2,3-difluorophenyl)-2-(4-{[7-(3-piperazin-1-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide;

10 *N*-(2,3-difluorophenyl)-2-{4-[(7-{3-[4-(2-hydroxyethyl)piperidin-1-yl]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;


N-(2,3-difluorophenyl)-2-{4-[(7-{3-[4-(hydroxymethyl)piperidin-1-yl]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;

N-(2,3-difluorophenyl)-2-{4-[(7-{3-[(2-hydroxyethyl)(isopropyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide;

15 2-{4-[(7-{3-[cyclopropyl(2-hydroxyethyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}-*N*-(2,3-difluorophenyl)acetamide;

or a salt, ester or prodrug thereof and more particularly a pharmaceutically acceptable salt thereof.


The present invention also provides a process for the preparation of a compound of
20 formula (I) or a salt, ester or prodrug thereof, which process comprises reacting a compound of formula (II)



(II)

where L is a suitable leaving group such as chloro, bromo, SMe etc.

25 with a compound of formula (III)




(III)

in the presence of hydrochloric acid in dioxane under an inert atmosphere, and thereafter if necessary:

- 5 i) converting a compound of the formula (I) into another compound of the formula (I); and/or
- ii) removing any protecting groups; and/or
- iii) forming a salt, ester or prodrug thereof.

The reaction is suitably effected in an organic solvent such as dimethyl acetamide or isopropanol at elevated temperatures of from 80°C to 120°C for 30 minutes to 2 hours.

- 10 The process may further comprise a process for the preparation of a compound of formula (II) when R^3 is $-X^3R^{13}$, which process comprises reacting a compound of formula (IV)



(IV)

- 15 with a compound of formula (V)




(V)

where L^1 is an appropriate leaving group such as chloro or L^1 is $-OH$ which is suitably activated by a reagent such as PPh_3 .

- 20 Compounds of formula (IV) and formula (V) are either known in the art or can be derived from other compounds known in the art by conventional methods which would be apparent to the skilled person from the literature. An analogous process exists for the preparation of a compound of formula (II) when R^3 is or is not $-X^3R^{13}$ and/or R^1 is $-X^1R^{11}$ and/or R^2 is $-X^2R^{12}$ and/or R^4 is $-X^4R^{14}$.

-29-


The process may further comprise a process for the preparation of a compound of formula (III) which process comprises the reaction of a compound of formula (VI)



(VI)

5

with a compound of formula (VII)



(VII)

The reaction is suitably effected in an organic solvent such as dimethylformamide or dimethylacetamide, with a base such as diisopropyl(ethyl)amine and with the addition of O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, maintaining a temperature of less than 40°C for 30 minutes to 2 hours.

Compounds of formula (VII) are known in the art or can be derived from other compounds known in the art by conventional methods which would be apparent to the skilled person from the literature.

A compound of formula (VI) when X is NR⁶, can be prepared by a process that comprises the:

- a) reaction of C₁₋₂₀alkyl azidoacetate with propiolic acid, followed by
 - b) reaction of the product of a) with a reagent such as diphenylphosphonyl azide.
- The reaction in a) is suitable effected in solvents such as chloroform, dichloromethane or toluene, at a temperature of 55°C to 100°C for 30 minutes to 5 hours, and the reaction in b) is effected in dioxane, under an inert atmosphere, under reflux for 2 to 7 hours.

Further provided is a process for the preparation of a compound of formula (IA) or a salt or ester thereof, which process comprises phosphorylation of a suitable compound of formula (I) by reacting a compound of formula (I) and tetrazole with di-*tert*-butyl diethylphosphoramidite in an appropriate organic solvent such as dimethylformamide or dimethylacetamide under an inert atmosphere, followed by (after 1 to 5 hours) the addition of hydrogen peroxide and sodium metabisulphite. Deprotection of the phosphate group then

-30-

yields a compound of formula (IA). Deprotection is suitably effected with hydrochloric acid in dioxane or dichloromethane (DCM) at ambient temperature for 6 to 30 hours.

Suitable reaction conditions are illustrated herein.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogen group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, *Protective Groups in Organic Synthesis*, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *tert*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such

-31-

as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *tert*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *tert*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound formula (I), or a pharmaceutically acceptable salt, ester or prodrug thereof, as defined herein in association with a pharmaceutically acceptable diluent or carrier.

Also provided is a pharmaceutical composition which comprises a compound of formula (IA), or a pharmaceutically acceptable salt or ester thereof, as defined herein in association with a pharmaceutically acceptable diluent or carrier.

The compositions of the invention may be in a form suitable for oral use (for example

-32-

as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

Suitable pharmaceutically acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc; preservative agents such as ethyl or propyl p-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal track, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, soya bean oil, coconut oil, or preferably olive oil, or any other acceptable vehicle.

Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxyethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or

-33-

condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and
5 hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl *p*-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

Oily suspensions may be formulated by suspending the active ingredient in a vegetable
10 oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

15 Dispersible or lyophilised powders and granules suitable for preparation of an aqueous suspension or solution by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients such as sweetening, flavouring and colouring agents,
20 may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum
25 tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, an esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.

30 Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable aqueous or oily suspension, solutions, emulsions or particular systems, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which have been mentioned above. A sterile injectable
5 preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in polyethylene glycol.

Suppository formulations may be prepared by mixing the active ingredient with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Suitable excipients
10 include, for example, cocoa butter and polyethylene glycols.

Topical formulations, such as creams, ointments, gels and aqueous or oily solutions or suspensions, may generally be obtained by formulating an active ingredient with a conventional, topically acceptable, vehicle or diluent using conventional procedure well known in the art.

15 Compositions for administration by insufflation may be in the form of a finely divided powder containing particles of average diameter of, for example, 30 μ m or much less preferably 5 μ m or less and more preferably between 5 μ m and 1 μ m, the powder itself comprising either active ingredient alone or diluted with one or more physiologically acceptable carriers such as lactose. The powder for insufflation is then conveniently retained
20 in a capsule containing, for example, 1 to 50mg of active ingredient for use with a turbo-inhaler device, such as is used for insufflation of the known agent sodium cromoglycate.

Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile
25 fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

For further information on formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

30 Therefore in a further aspect of the invention there is provided a compound of formula (I), or a pharmaceutically acceptable salt, ester or prodrug thereof, for use in therapy. In addition a compound of formula (IA) or a pharmaceutically acceptable salt or ester thereof is

provided for use in therapy.

Further provided is a compound of formula (I), or a pharmaceutically acceptable salt, ester or prodrug thereof, for use as a medicament and also provided is a compound of formula (IA), or a pharmaceutically acceptable salt or ester thereof, for use as a medicament. Another aspect of the invention provides a compound of formula (I), or a pharmaceutically acceptable salt, ester or prodrug thereof, for use as a medicament for the treatment of hyperproliferative diseases such as cancer and in particular colorectal, breast, lung, prostate, bladder, renal or pancreatic cancer or leukaemia or lymphoma. Also provided is a compound of formula (IA), or a pharmaceutically acceptable salt or ester thereof, for use as a medicament for the treatment of hyperproliferative diseases such as cancer and in particular colorectal, breast, lung, prostate, bladder, renal or pancreatic cancer or leukaemia or lymphoma.

Additionally a compound of formula (I), or a pharmaceutically acceptable salt, ester or prodrug thereof is provided for use in a method of treatment of a warm-blooded animal such as man by therapy. A compound of formula (IA) or a pharmaceutically acceptable salt or ester thereof is also provided for use in a method of treatment of a warm-blooded animal such as man by therapy. Another aspect of the invention provides a compound of formula (I), or a pharmaceutically acceptable salt, ester or prodrug thereof, for use in a method of treatment of hyperproliferative diseases such as cancer and in particular colorectal, breast, lung, prostate, bladder, renal or pancreatic cancer or leukaemia or lymphoma. Also provided is a compound of formula (IA), or a pharmaceutically acceptable salt or ester thereof, for use in a method of treatment of hyperproliferative diseases such as cancer and in particular colorectal, breast, lung, prostate, bladder, renal or pancreatic cancer or leukaemia or lymphoma.

In another aspect of the invention, there is provided the use of a compound of formula (I) or a pharmaceutically acceptable salt, ester or prodrug thereof, in the preparation of a medicament for the treatment of a disease where the inhibition of one or more Aurora kinase(s) is beneficial. The use of a compound of formula (IA) or a pharmaceutically acceptable salt or ester thereof in the preparation of a medicament for the treatment of a disease where the inhibition of one or more Aurora kinase(s) is beneficial is also provided. In particular it is envisaged that inhibition of Aurora-A kinase and/or Aurora-B kinase may be beneficial. Preferably inhibition of Aurora-B kinase is beneficial. In another aspect of the invention, there is provided the use of a compound of formula (I) or a pharmaceutically acceptable salt, ester or prodrug thereof, in the preparation of a medicament for the treatment of hyperproliferative diseases such as cancer and in particular colorectal, breast, lung,

prostate, bladder, renal or pancreatic cancer or leukaemia or lymphoma. Also provided is the use of a compound of formula (IA) or a pharmaceutically acceptable salt or ester thereof in the preparation of a medicament for the treatment of hyperproliferative diseases such as cancer and in particular colorectal, breast, lung, prostate, bladder, renal or pancreatic cancer or
5 leukaemia or lymphoma.

According to yet another aspect, there is provided a compound of formula (I) or a pharmaceutically acceptable salt, ester or prodrug thereof for use in the method of treating a human suffering from a disease in which the inhibition of one or more Aurora kinases is beneficial, comprising the steps of administering to a person in need thereof a therapeutically
10 effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, ester or prodrug thereof. Further provided is a compound of formula (IA) or a pharmaceutically acceptable salt thereof for use in the method of treating a human suffering from a disease in which the inhibition of one or more Aurora kinases is beneficial, comprising the steps of administering to a person in need thereof a therapeutically effective amount of a compound of
15 formula (IA) or a pharmaceutically acceptable salt thereof. In particular it is envisaged that inhibition of Aurora-A kinase and/or Aurora-B kinase may be beneficial. Preferably inhibition of Aurora-B kinase is beneficial. Further provided is a compound of formula (I) or a pharmaceutically acceptable salt, ester or prodrug thereof for use in the method of treating a human suffering from a hyperproliferative disease such as cancer and in particular colorectal,
20 breast, lung, prostate, bladder, renal or pancreatic cancer or leukaemia or lymphoma, comprising the steps of administering to a person in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, ester or prodrug thereof. A compound of formula (IA) is also provided for use in the method of treating a human suffering from a hyperproliferative disease such as cancer and in particular colorectal,
25 breast, lung, prostate, bladder, renal or pancreatic cancer or leukaemia or lymphoma, comprising the steps of administering to a person in need thereof a therapeutically effective amount of a compound of formula (IA) or a pharmaceutically acceptable salt or ester thereof. The use of a compound of formula (I) or a pharmaceutically acceptable salt, ester or prodrug thereof in any of the methods of treating a human described above also form aspects of this
30 invention. Additionally the use of a compound of formula (IA) or a pharmaceutically acceptable salt or ester thereof in any of the methods of treating a human described above form other aspects of this invention.

For the above mentioned therapeutic uses the dose administered will vary with the

compound employed, the mode of administration, the treatment desired, the disorder indicated and the age and sex of the animal or patient. The size of the dose would thus be calculated according to well known principles of medicine.

In using a compound of formula (I) or formula (IA) for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.05 mg/kg to 50 mg/kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.05 mg/kg to 25 mg/kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.05 mg/kg to 25 mg/kg body weight will be used.

The treatment defined herein may be applied as a sole therapy or may involve, in addition to the compound of the invention, conventional surgery or radiotherapy or chemotherapy. Such chemotherapy may include one or more of the following categories of anti-tumour agents :-

- 15 (i) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as alkylating agents (for example cis-platin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan and nitrosoureas); antimetabolites (for example antifolates such as fluoropyrimidines like 5-fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside and hydroxyurea; antitumour antibiotics (for example 20 anthracyclines like adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine and vinorelbine and taxoids like taxol and taxotere); and topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan and camptothecin);
- 25 (ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene and idoxifene), oestrogen receptor down regulators (for example fulvestrant), antiandrogens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example goserelin, leuprorelin and buserelin), progestogens (for example megestrol acetate), aromatase inhibitors (for 30 example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 α -reductase such as finasteride;

- (iii) Agents which inhibit cancer cell invasion (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function);
- (iv) inhibitors of growth factor function, for example such inhibitors include growth factor antibodies, growth factor receptor antibodies (for example the anti-erb2 antibody
- 5 trastuzumab [Herceptin™] and the anti-erb1 antibody cetuximab [C225]), farnesyl transferase inhibitors, tyrosine kinase inhibitors and serine-threonine kinase inhibitors, for example inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors such as *N*-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, AZD1839), *N*-(3-ethynylphenyl)-6,7-
- 10 bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-*N*-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)quinazolin-4-amine (CI 1033)), for example inhibitors of the platelet-derived growth factor family and for example inhibitors of the hepatocyte growth factor family;
- (v) antiangiogenic agents such as those which inhibit the effects of vascular endothelial
- 15 growth factor, (for example the anti-vascular endothelial cell growth factor antibody bevacizumab [Avastin™], compounds such as those disclosed in International Patent Applications WO 97/22596, WO 97/30035, WO 97/32856 and WO 98/13354) and compounds that work by other mechanisms (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function and angiostatin);
- 20 (vi) vascular damaging agents such as Combretastatin A4 and compounds disclosed in International Patent Applications WO 99/02166, WO00/40529, WO 00/41669, WO01/92224, WO02/04434 and WO02/08213;
- (vii) antisense therapies, for example those which are directed to the targets listed above, such as ISIS 2503, an anti-ras antisense;
- 25 (viii) gene therapy approaches, including for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; and
- 30 (ix) immunotherapy approaches, including for example ex-vivo and *in vivo* approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor,

-39-

approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines and approaches using anti-idiotypic antibodies.

In addition a compound of the invention or a pharmaceutically acceptable salt, ester or prodrug thereof, may be used in combination with one or more cell cycle inhibitors. In particular with cell cycle inhibitors which inhibit bub1, bubR1 or CDK.

Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate dosing of the individual components of the treatment. Such combination products employ the compounds of this invention within the dosage range described herein and the other pharmaceutically-active agent within its approved dosage range.

In addition to their use in therapeutic medicine, a compound of formula (I) and a pharmaceutically acceptable salt, ester or prodrug thereof are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of cell cycle activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

The compounds of the invention inhibit the serine-threonine kinase activity of the Aurora kinases, in particular Aurora-A kinase and/or Aurora-B kinase and thus inhibit the cell cycle and cell proliferation. Compounds which inhibit Aurora-B kinase are of particular interest. These properties may be assessed for example, using one or more of the procedures set out below.

(a) In Vitro Aurora-A kinase inhibition test

This assay determines the ability of a test compound to inhibit serine-threonine kinase activity. DNA encoding Aurora-A may be obtained by total gene synthesis or by cloning. This DNA may then be expressed in a suitable expression system to obtain polypeptide with serine-threonine kinase activity. In the case of Aurora-A, the coding sequence was isolated from cDNA by polymerase chain reaction (PCR) and cloned into the BamH1 and Not1 restriction endonuclease sites of the baculovirus expression vector pFastBac HTc (GibcoBRL/Life technologies). The 5' PCR primer contained a recognition sequence for the restriction endonuclease BamH1 5' to the Aurora-A coding sequence. This allowed the insertion of the

Aurora-A gene in frame with the 6 histidine residues, spacer region and rTEV protease cleavage site encoded by the pFastBac HTc vector. The 3' PCR primer replaced the Aurora-A stop codon with additional coding sequence followed by a stop codon and a recognition sequence for the restriction endonuclease NotI. This additional coding sequence (5' TAC CCA TAC GAT GTT CCA GAT TAC GCT TCT TAA 3') encoded for the polypeptide sequence YPYDVDPDYAS. This sequence, derived from the influenza hemagglutinin protein, is frequently used as a tag epitope sequence that can be identified using specific monoclonal antibodies. The recombinant pFastBac vector therefore encoded for an N-terminally 6 his tagged, C terminally influenza hemagglutinin epitope tagged Aurora-A protein. Details of the methods for the assembly of recombinant DNA molecules can be found in standard texts, for example Sambrook et al. 1989, Molecular Cloning - A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory press and Ausubel et al. 1999, Current Protocols in Molecular Biology, John Wiley and Sons Inc.

Production of recombinant virus can be performed following manufacturer's protocol from GibcoBRL. Briefly, the pFastBac-1 vector carrying the Aurora-A gene was transformed into E. coli DH10Bac cells containing the baculovirus genome (bacmid DNA) and via a transposition event in the cells, a region of the pFastBac vector containing gentamycin resistance gene and the Aurora-A gene including the baculovirus polyhedrin promoter was transposed directly into the bacmid DNA. By selection on gentamycin, kanamycin, tetracycline and X-gal, resultant white colonies should contain recombinant bacmid DNA encoding Aurora-A. Bacmid DNA was extracted from a small scale culture of several BH10Bac white colonies and transfected into *Spodoptera frugiperda* Sf21 cells grown in TC100 medium (GibcoBRL) containing 10% serum using CellFECTIN reagent (GibcoBRL) following manufacturer's instructions. Virus particles were harvested by collecting cell culture medium 72 hrs post transfection. 0.5 mls of medium was used to infect 100 ml suspension culture of Sf21s containing 1×10^7 cells/ml. Cell culture medium was harvested 48 hrs post infection and virus titre determined using a standard plaque assay procedure. Virus stocks were used to infect Sf9 and "High 5" cells at a multiplicity of infection (MOI) of 3 to ascertain expression of recombinant Aurora-A protein.

For the large scale expression of Aurora-A kinase activity, Sf21 insect cells were grown at 28°C in TC100 medium supplemented with 10% foetal calf serum (Viralex) and 0.2% F68 Pluronic (Sigma) on a Wheaton roller rig at 3 r.p.m. When the cell density reached

1.2x10⁶ cells ml⁻¹ they were infected with plaque-pure Aurora-A recombinant virus at a multiplicity of infection of 1 and harvested 48 hours later. All subsequent purification steps were performed at 4°C. Frozen insect cell pellets containing a total of 2.0 x 10⁸ cells were thawed and diluted with lysis buffer (25 mM HEPES (N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulphonic acid]) pH7.4 at 4°C, 100 mM KCl, 25 mM NaF, 1 mM Na₃VO₄, 1 mM PMSF (phenylmethylsulphonyl fluoride), 2 mM 2-mercaptoethanol, 2 mM imidazole, 1 µg/ml aprotinin, 1 µg/ml pepstatin, 1 µg/ml leupeptin), using 1.0 ml per 3 x 10⁷ cells. Lysis was achieved using a dounce homogeniser, following which the lysate was centrifuged at 41,000g for 35 minutes. Aspirated supernatant was pumped onto a 5 mm diameter chromatography column containing 500 µl Ni NTA (nitrilo-tri-acetic acid) agarose (Qiagen, product no. 30250) which had been equilibrated in lysis buffer. A baseline level of UV absorbance for the eluent was reached after washing the column with 12 ml of lysis buffer followed by 7 ml of wash buffer (25 mM HEPES pH7.4 at 4°C, 100 mM KCl, 20 mM imidazole, 2 mM 2-mercaptoethanol). Bound Aurora-A protein was eluted from the column using elution buffer (25 mM HEPES pH7.4 at 4°C, 100 mM KCl, 400 mM imidazole, 2 mM 2-mercaptoethanol). An elution fraction (2.5 ml) corresponding to the peak in UV absorbance was collected. The elution fraction, containing active Aurora-A kinase, was dialysed exhaustively against dialysis buffer (25 mM HEPES pH7.4 at 4°C, 45% glycerol (v/v), 100 mM KCl, 0.25% Nonidet P40 (v/v), 1 mM dithiothreitol).

Each new batch of Aurora-A enzyme was titrated in the assay by dilution with enzyme diluent (25mM Tris-HCl pH7.5, 12.5mM KCl, 0.6mM DTT). For a typical batch, stock enzyme is diluted 1 in 666 with enzyme diluent & 20µl of dilute enzyme is used for each assay well. Test compounds (at 10mM in dimethylsulphoxide (DMSO)) were diluted with water & 10µl of diluted compound was transferred to wells in the assay plates. "Total" & "blank" control wells contained 2.5% DMSO instead of compound. Twenty microlitres of freshly diluted enzyme was added to all wells, apart from "blank" wells. Twenty microlitres of enzyme diluent was added to "blank" wells. Twenty microlitres of reaction mix (25mM Tris-HCl, 78.4mM KCl, 2.5mM NaF, 0.6mM dithiothreitol, 6.25mM MnCl₂, 6.25mM ATP, 7.5µM peptide substrate [biotin-LRRWSLGLRRWSLGLRRWSLGLRRWSLGL]) containing 0.2µCi [³³P]ATP (Amersham Pharmacia, specific activity ≥2500Ci/mmol) was then added to all test wells to start the reaction. The plates were incubated at room temperature for 60 minutes. To stop the reaction 100µl 20% v/v orthophosphoric acid was added to all wells. The

peptide substrate was captured on positively-charged nitrocellulose P30 filtermat (Whatman) using a 96-well plate harvester (TomTek) & then assayed for incorporation of ^{33}P with a Beta plate counter. "Blank" (no enzyme) and "total" (no compound) control values were used to determine the dilution range of test compound which gave 50% inhibition of enzyme activity.

- 5 In this test, the compounds of the invention generally give 50% inhibition of enzyme activity at concentrations of 1nM to 1000nM and in particular compound 1 in Table 1 gave 50% inhibition of enzyme activity at a concentration of 0.9 μM and compound 4 in Table 2 gave 50% inhibition of enzyme activity at a concentration of 0.5 μM

10 (b) In Vitro Aurora-B kinase inhibition test

This assay determines the ability of a test compound to inhibit serine-threonine kinase activity. DNA encoding Aurora-B may be obtained by total gene synthesis or by cloning. This DNA may then be expressed in a suitable expression system to obtain polypeptide with serine-threonine kinase activity. In the case of Aurora-B, the coding sequence was isolated from
15 cDNA by polymerase chain reaction (PCR) and cloned into the pFastBac system in a manner similar to that described above for Aurora-A (i.e. to direct expression of a 6-histidine tagged Aurora-B protein).

For the large scale expression of Aurora-B kinase activity, Sf21 insect cells were grown at 28°C in TC100 medium supplemented with 10% foetal calf serum (Viralex) and
20 0.2% F68 Pluronic (Sigma) on a Wheaton roller rig at 3 r.p.m. When the cell density reached 1.2×10^6 cells ml^{-1} they were infected with plaque-pure Aurora-B recombinant virus at a multiplicity of infection of 1 and harvested 48 hours later. All subsequent purification steps were performed at 4°C. Frozen insect cell pellets containing a total of 2.0×10^8 cells were thawed and diluted with lysis buffer (50 mM HEPES (N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulphonic acid]) pH7.5 at 4°C, 1 mM Na_3VO_4 , 1 mM PMSF (phenylmethylsulphonyl
25 fluoride), 1 mM dithiothreitol, 1 $\mu\text{g}/\text{ml}$ aprotinin, 1 $\mu\text{g}/\text{ml}$ pepstatin, 1 $\mu\text{g}/\text{ml}$ leupeptin), using 1.0 ml per 2×10^7 cells. Lysis was achieved using a sonication homogeniser, following which the lysate was centrifuged at 41,000g for 35 minutes. Aspirated supernatant was pumped onto a 5 mm diameter chromatography column containing 1.0 ml CM sepharose Fast Flow
30 (Amersham Pharmacia Biotech) which had been equilibrated in lysis buffer. A baseline level of UV absorbance for the eluent was reached after washing the column with 12 ml of lysis buffer followed by 7 ml of wash buffer (50 mM HEPES pH7.4 at 4°C, 1 mM dithiothreitol).

-43-

Bound Aurora-B B protein was eluted from the column using a gradient of elution buffer (50 mM HEPES pH7.4 at 4°C, 0.6 M NaCl, 1 mM dithiothreitol, running from 0% elution buffer to 100% elution buffer over 15 minutes at a flowrate of 0.5 ml/min). Elution fractions (1.0 ml) corresponding to the peak in UV absorbance was collected. Elution fractions were dialysed exhaustively against dialysis buffer (25 mM HEPES pH7.4 at 4°C, 45% glycerol (v/v), 100 mM KCl, 0.05% (v/v) IGEPAL CA630 (Sigma Aldrich), 1 mM dithiothreitol). Dialysed fractions were assayed for Aurora-B kinase activity.

Each new batch of Aurora-B enzyme was titrated in the assay by dilution with enzyme diluent (25mM Tris-HCl pH7.5, 12.5mM KCl, 0.6mM DTT). For a typical batch, stock enzyme is diluted 1 in 40 with enzyme diluent & 20µl of dilute enzyme is used for each assay well. Test compounds (at 10mM in dimethylsulphoxide (DMSO)) were diluted with water & 10µl of diluted compound was transferred to wells in the assay plates. "Total" & "blank" control wells contained 2.5% DMSO instead of compound. Twenty microlitres of freshly diluted enzyme was added to all wells, apart from "blank" wells. Twenty microlitres of enzyme diluent was added to "blank" wells. Twenty microlitres of reaction mix (25mM Tris-HCl, 78.4mM KCl, 2.5mM NaF, 0.6mM dithiothreitol, 6.25mM MnCl₂, 37.5mM ATP, 25µM peptide substrate [biotin-LRRWSLGLRRWSLGLRRWSLGLRRWSLG]) containing 0.2µCi [³³P]ATP (Amersham Pharmacia, specific activity ≥2500Ci/mmol) was then added to all test wells to start the reaction. The plates were incubated at room temperature for 60 minutes. To stop the reaction 100µl 20% v/v orthophosphoric acid was added to all wells. The peptide substrate was captured on positively-charged nitrocellulose P30 filtermat (Whatman) using a 96-well plate harvester (TomTek) & then assayed for incorporation of ³³P with a Beta plate counter. "Blank" (no enzyme) and "total" (no compound) control values were used to determine the dilution range of test compound which gave 50% inhibition of enzyme activity. In this test, the compounds of the invention generally give 50% inhibition of enzyme activity at concentrations of 1nM to 1000nM and in particular compound 1 in Table 1 gave 50% inhibition of enzyme activity at a concentration of 0.1µM and compound 4 in Table 2 gave 50% inhibition of enzyme activity at a concentration of 0.1µM.

30 (c) In Vitro cell proliferation assay

This and other assays can be used to determine the ability of a test compound to inhibit the growth of adherent mammalian cell lines, for example the human tumour cell line SW620

-44-

(ATCC CCL-227). This assay determines the ability of a test compound to inhibit the incorporation of the thymidine analogue, 5'-bromo-2'-deoxy-uridine (BrdU) into cellular DNA. SW620 or other adherent cells were typically seeded at 1×10^5 cells per well in L-15 media (GIBCO) plus 5% foetal calf serum, 1% L-glutamine (100 μ l / well) in 96 well tissue culture treated 96 well plates (Costar) and allowed to adhere overnight. The following day the cells were dosed with compound (diluted from 10mM stock in DMSO using L-15 (with 5% FCS, 1% L-glutamine). Untreated control wells and wells containing a compound known to give 100% inhibition of BrdU incorporation were included on each plate. After 48 hours in the presence / absence of test compound the ability of the cells to incorporate BrdU over a 2 hour labelling period was determined using a Boehringer (Roche) Cell Proliferation BrdU ELISA kit (cat. No. 1 647 229) according to manufacturers directions. Briefly, 15 μ l of BrdU labelling reagent (diluted 1:100 in media – L-15, 5% FCS, 1% L-glutamine) was added to each well and the plate returned to a humidified (+5% CO₂) 37°C incubator for 2 hours. After 2 hours the labelling reagent was removed by decanting and tapping the plate on a paper towel. FixDenat solution (50 μ l per well) was added and the plates incubated at room temperature for 45mins with shaking. The FixDenat solution was removed by decanting and tapping the inverted plate on a paper towel. The plate was then washed once with phosphate buffered saline (PBS) and 100 μ l /well of Anti-BrdU-POD antibody solution (diluted 1:100 in antibody dilution buffer) added. The plate was then incubated at room temperature with shaking for 90min. Unbound Anti-BrdU-POD antibody was removed by decanting and washing the plate 4 times with PBS before being blotted dry. TMB substrate solution was added (100 μ l/well) and incubated for approximately 10 minutes at room temperature with shaking until a colour change was apparent. The optical density of the wells was then determined at 690nm wavelength using a Titertek Multiscan plate reader. The values from compound treated, untreated and 100% inhibition controls were used to determine the dilution range of a test compound that gave 50% inhibition of BrdU incorporation. The compounds of the invention are generally active at 1nM to 100 μ M in this test.

(d) In Vitro cell cycle analysis assay

This assay determines the ability of a test compound to arrest cells in specific phases of the cell cycle. Many different mammalian cell lines could be used in this assay and SW620 cells are included here as an example. SW620 cells were seeded at 7×10^5 cells per T25 flask

-45-

(Costar) in 5 ml L-15 (5% FCS, 1% L-glutamine). Flasks were then incubated overnight in a humidified 37°C incubator with 5% CO₂. The following day, 5µl of L-15 (5% FCS, 1% L-glutamine) carrying the appropriate concentration of test compound solubilised in DMSO was added to the flask. A no compound control treatment was also included (0.5% DMSO). The cells were then incubated for a defined time (24 hours) with compound. After this time the media was aspirated from the cells and they were washed with 5ml of prewarmed (37°C) sterile PBSA, then detached from the flask by brief incubation with trypsin and followed by resuspension in 5ml of 1% Bovine Serum Albumin (BSA, Sigma-Aldrich Co.) in sterile PBSA. The samples were then centrifuged at 2200rpm for 10 min. The supernatant was aspirated to leave 200µl of the PBS/BSA solution. The pellet was resuspended in this 200µl of solution by pipetting 10 times to create a single cell suspension. One ml of ice-cold 80% ethanol was slowly added to each cell suspension and the samples stored at -20°C overnight or until required for staining. Cells were pelleted by centrifugation, ethanol aspirated off and pellets resuspended in 200µl PBS containing 100µg/ml RNase (Sigma Aldrich) & 10µg/ml Propidium Iodide (Sigma Aldrich). Cell suspensions were incubated at 37°C for 30min, a further 200µl PBS added and samples stored in the dark at 4°C overnight.

Each sample was then syringed 10 times using 21-guage needle. The samples were then transferred to LPS tubes and DNA content per cell analysed by Fluorescence activated cell sorting (FACS) using a FACScan flow cytometer (Becton Dickinson). Typically 30,000 events were counted and recorded using CellQuest v1.1 software (Verity Software). Cell cycle distribution of the population was calculated using Modfit software (Verity Software) and expressed as percentage of cells with 2N (G0/G1), 2N-4N (S phase) and with 4N (G2/M) DNA content.

The compounds of the invention are generally active in this test at 1nM to 10µM.

The invention will now be illustrated in the following examples, in which standard techniques known to the skilled chemist and techniques analogous to those described in these examples may be used where appropriate, and in which, unless otherwise stated:

- (i) evaporations were carried out by rotary evaporation *in vacuo* and work up procedures were carried out after removal of residual solids such as drying agents by filtration;
- (ii) operations were carried out at ambient temperature, typically in the range 18-25°C and in air unless stated, or unless the skilled person would otherwise operate under an atmosphere of

an inert gas such as argon;

(iii) column chromatography (by the flash procedure) and medium pressure liquid chromatography (MPLC) were performed on Merck Kieselgel silica (Art. 9385);

(iv) yields are given for illustration only and are not necessarily the maximum attainable;

5 (v) the structures of the end products of the formula (I) were generally confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic resonance chemical shift values were measured in deuterated dimethyl sulphoxide (DMSO d_6) (unless otherwise stated) on the delta scale (ppm downfield from tetramethylsilane) using one of the following four instruments

- 10
- Varian Gemini 2000 spectrometer operating at a field strength of 300 MHz
 - Bruker DPX300 spectrometer operating at a field strength of 300MHz
 - JEOL EX 400 spectrometer operating at a field strength of 400 MHz
 - Bruker Avance 500 spectrometer operating at a field strength of 500MHz

Peak multiplicities are shown as follows: s, singlet; d, doublet; dd, double doublet; t, triplet; q, quartet; qu, quintet; m, multiplet; br s, broad singlet;

15 (vi) robotic synthesis was carried out using a Zymate XP robot, with solution additions via a Zymate Master Laboratory Station and stirred via a Stem RS5000 Reacto-Station at 25°C;

(vii) work up and purification of reaction mixtures from robotic synthesis was carried out as follows: evaporations were carried out *in vacuo* using a Genevac HT 4; column

20 chromatography was performed using either an Anachem Sympur MPLC system on silica using 27 mm diameter columns filled with Merck silica (60 μ m, 25 g); the structures of the final products were confirmed by LCMS (liquid chromatography mass spectrometry) on a Waters 2890 / ZMD micromass system using the following and are quoted as retention time (RT) in minutes:

25 Column: waters symmetry C18 3.5 μ m 4.6x50 mm

Solvent A: H₂O

Solvent B: CH₃CN

Solvent C : MeOH + 5% HCOOH

Flow rate: 2.5 ml / min

30 Run time: 5 minutes with a 4.5 minute gradient from 0-100% C

Wavelength: 254 nm, bandwidth 10 nm

Mass detector: ZMD micromass

Injection volume 0.005 ml

(viii) Analytical LCMS for compounds which had not been prepared by robotic synthesis was performed on a Waters Alliance HT system using the following and are quoted as retention time (RT) in minutes:

Column: 2.0 mm x 5 cm Phenomenex Max-RP 80A
5 Solvent A: Water
Solvent B: Acetonitrile
Solvent C: Methanol / 1% formic acid or Water / 1% formic acid
Flow rate: 1.1 ml / min
Run time: 5 minutes with a 4.5 minute gradient from 0-95% B + constant 5%
10 solvent C
Wavelength: 254 nm, bandwidth 10 nm
Injection volume 0.005 ml
Mass detector: Micromass ZMD

(ix) Preparative high performance liquid chromatography (HPLC) was performed on either
15 - Waters preparative LCMS instrument, with retention time (RT) measured in minutes:

Column: β -basic Hypercil (21x100 mm) 5 μ m
Solvent A: Water / 0.1% Ammonium carbonate
Solvent B: Acetonitrile
Flow rate: 25 ml / min
20 Run time: 10 minutes with a 7.5 minute gradient from 0-100% B
Wavelength: 254 nm, bandwidth 10 nm
Injection volume 1 - 1.5 ml
Mass detector : Micromass ZMD


- Gilson preparative HPLC instrument, with retention time (RT) measured in minutes:

25 Column: 21 mm x 15 cm Phenomenex Luna2 C18
Solvent A: Water + 0.1% trifluoroacetic acid,
Solvent B: Acetonitrile + 0.1% trifluoroacetic acid
Flow rate: 21 ml / min
Run time: 20 minutes with various 10 minute gradients from 5-100% B
30 Wavelength: 254 nm, bandwidth 10 nm
Injection volume 0.1-4.0 ml

(x) intermediates were not generally fully characterised and purity was assessed by thin layer


-48-

chromatography (TLC), HPLC, infra-red (IR), MS or NMR analysis.

Table 1

5

Compound	R ²	R ⁵
1	OMe	3-fluorophenyl
2	H	3-fluorophenyl
3	H	2,3-difluorophenyl

Table 2

10

Compound	R ²	R	R ³
4	OMe	H	3-[(2-hydroxyethyl)(propyl)amino]propoxy
5	OMe	H	3-[(2 <i>S</i>)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy
6	H	H	3-[(2-hydroxyethyl)(propyl)amino]propoxy
7	H	H	3-[(2 <i>S</i>)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy
8	H	H	3-morpholin-4-ylpropoxy
9	H	H	3-piperidin-1-ylpropoxy

-49-

10	H	H	3-pyrrolidin-1-ylpropoxy
11	H	H	3-[(2-hydroxy-1,1-dimethylethyl)amino]propoxy
12	H	H	3-(cyclopropylamino)propoxy
13	H	H	3-[[2-(dimethylamino)ethyl](methyl)amino]propoxy
14	H	H	3-(4-methylpiperazin-1-yl)propoxy
15	H	H	3-[(2R)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy
16	H	H	3-(4-hydroxypiperidin-1-yl)propoxy
17	H	H	3-[ethyl(2-hydroxyethyl)amino]propoxy
18	H	H	3-[4-(2-hydroxyethyl)piperazin-1-yl]propoxy
19	H	H	3-piperazin-1-ylpropoxy
20	H	H	3-[4-(2-hydroxyethyl)piperidin-1-yl]propoxy
21	H	H	3-[4-(hydroxymethyl)piperidin-1-yl]propoxy
22	H	H	3-[(2-hydroxyethyl)(isopropyl)amino]propoxy
23	H	H	3-[cyclopropyl(2-hydroxyethyl)amino]propoxy
24	H	F	3-morpholin-4-ylpropoxy
25	H	F	3-piperidin-1-ylpropoxy
26	H	F	3-pyrrolidin-1-ylpropoxy
27	H	F	3-[(2-hydroxy-1,1-dimethylethyl)amino]propoxy
28	H	F	3-(cyclopropylamino)propoxy
29	H	F	3-[[2-(dimethylamino)ethyl](methyl)amino]propoxy
30	H	F	3-(4-methylpiperazin-1-yl)propoxy
31	H	F	3-[(2R)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy
32	H	F	3-(4-hydroxypiperidin-1-yl)propoxy
33	H	F	3-[ethyl(2-hydroxyethyl)amino]propoxy
34	H	F	3-[4-(2-hydroxyethyl)piperazin-1-yl]propoxy
35	H	F	3-piperazin-1-ylpropoxy
36	H	F	3-[4-(2-hydroxyethyl)piperidin-1-yl]propoxy

-50-

37	H	F	3-[4-(hydroxymethyl)piperidin-1-yl]propoxy
38	H	F	3-[(2-hydroxyethyl)(isopropyl)amino]propoxy
39	H	F	3-[cyclopropyl(2-hydroxyethyl)amino]propoxy

Example 1 - Preparation of compound 1 in table 1 - 2-(4-[[7-(3-chloropropoxy)-6-methoxyquinazolin-4-yl]amino]-1H-1,2,3-triazol-1-yl)-N-(3-fluorophenyl)acetamide

2-(4-amino-1H-1,2,3-triazol-1-yl)-N-(3-fluorophenyl)acetamide (400 mg, 1.7 mmol) was added to a solution of 4-chloro-7-(3-chloropropoxy)-6-methoxyquinazoline (488 mg, 1.7 mmol) in dimethyl acetamide (15 ml). A solution of hydrochloric acid in dioxane (4.0 N, 235 μ l, 1.7 mmol) was added to the reaction mixture and the resulting solution was heated at 90 °C for 50 minutes causing a dense precipitate to form. The reaction mixture was cooled and diluted with isopropanol. The solid was recovered by suction filtration, washed with ethyl acetate and dried *in vacuo* to give compound 1 in table 1 (860 mg, 85 % yield) :

¹H-NMR (DMSO d₆) : 9.05 (s, 1H), 8.72 (s, 1H), 8.37 (s, 1H), 7.61 (m, 1H), 7.43 (s, 1H), 7.36 (m, 2H), 6.93 (t, 1H), 5.51 (s, 2H), 4.35 (t, 2H), 4.04 (s, 3H), 3.85 (t, 2H), 2.33 (m, 2H) : MS (+ve ESI): 486.1 (M+H)⁺.

2-(4-amino-1H-1,2,3-triazol-1-yl)-N-(3-fluorophenyl)acetamide, used as starting material, was obtained as follows:

a) Ethyl azidoacetate (3.96 ml of a 3.26 N solution in dichloromethane, 10 mmol) was added to a solution of propionic acid (700 mg, 10 mmol) in toluene (5 ml) and the reaction heated at reflux for 1 hour. The reaction was cooled and the solid was recovered, washed with diethyl ether and dried *in vacuo* to give 1-(2-ethoxy-2-oxoethyl)-1H-1,2,3-triazole-4-carboxylic acid (1.4 g, 70 % yield) :

¹H-NMR (DMSO d₆) : 8.67 (s, 1H), 5.46 (s, 2H), 4.19 (q, 2H), 1.23 (t, 3H) : MS (+ve ESI): 200.2 (M+H)⁺.

b) Diphenylphosphoryl azide (11.7 g, 42 mmol) was slowly added to a suspension of 1-(2-ethoxy-2-oxoethyl)-1H-1,2,3-triazole-4-carboxylic acid (7.56 g, 38 mmol) in a mixture of dry dioxane (100 ml) and 2-methylpropan-2-ol (50 ml) under argon. The solution was slowly heated to reflux and heated at reflux for 5 hours. The reaction mixture was cooled, concentrated *in vacuo*, and the residual oil diluted with a mixture of ethyl acetate (100 ml) and diethyl ether (50 ml). The solution was washed with water and brine before being concentrated *in vacuo*. Purification by chromatography on silica gel, eluting with

-51-

dichloromethane : ethyl acetate (9:1 to 7:3) gave ethyl {4-[(*tert*-butoxycarbonyl)amino]-1*H*-1,2,3-triazol-1-yl}acetate as a white solid (5.52 g, 54 % yield) :

¹H-NMR (DMSO *d*₆): 10.05 (s, 1H), 7.94 (s, 1H), 5.31 (s, 2H), 4.17 (q, 2H), 1.46 (s, 9H), 1.22 (t, 3H) :

5 MS (+ve ESI) : 271.3 (M+H)⁺.

c) A solution of ethyl {4-[(*tert*-butoxycarbonyl)amino]-1*H*-1,2,3-triazol-1-yl}acetate (2.7 g, 10 mmol) in ethanol (54 ml) and 2.0 N aqueous sodium hydroxide (10 ml, 20 mmol) was stirred at ambient temperature for 3 hours. The pH of the solution was then adjusted to 7, the solvent was evaporated *in vacuo*, and the pH was adjusted to 3. The precipitate was collected
10 by suction filtration, washed with water and dried to give {4-[(*tert*-butoxycarbonyl)amino]-1*H*-1,2,3-triazol-1-yl}acetic acid (2.35 g, 97 % yield) :

¹H-NMR (DMSO *d*₆) : 10.03 (s, 1H), 7.91 (s, 1H), 5.19 (s, 2H), 1.46 (s, 9H) :

MS (+ve ESI): 243.2 (M+H)⁺.

d) 3-Fluoroaniline (670 mg, 6 mmol) was added to a solution of {4-[(*tert*-
15 butoxycarbonyl)amino]-1*H*-1,2,3-triazol-1-yl}acetic acid (1.21 g, 5 mmol) in dimethyl formamide (12 ml) and diisopropylethylamine (770 mg, 6 mmol).
O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (2.08 g, 5.5 mmol) was added to the solution at such a rate to keep the temperature of the reaction medium below 30 °C. The mixture was stirred for 40 minutes, diluted with ethyl acetate (40 ml) and
20 diethyl ether (40 ml) and then washed with i) sodium bicarbonate solution, ii) 0.5 N hydrochloric acid and iii) brine. The organic phase was concentrated *in vacuo* to give *tert*-butyl (1-{2-[(3-fluorophenyl)amino]-2-oxoethyl}-1*H*-1,2,3-triazol-4-yl)carbamate (1.38 g, 82 % yield) :

¹H-NMR (DMSO *d*₆) : 10.65 (s, 1H), 10.04 (s, 1H), 7.95 (m, 1H), 7.55 (m, 1H), 7.38 (m, 1H),
25 7.30 (d, 1H), 6.93 (m, 1H), 5.28 (s, 1H), 1.46 (s, 9H) :

MS (+ve ESI): 336.2 (M+H)⁺.

e) Trifluoroacetic acid (6 ml) was added to a suspension of *tert*-butyl (1-{2-[(3-fluorophenyl)amino]-2-oxoethyl}-1*H*-1,2,3-triazol-4-yl)carbamate (1.5 g, 4.5 mmol) in dichloromethane (12 ml), and the reaction was stirred at 45 °C for 1.5 hours. The solvents
30 were evaporated *in vacuo* and aqueous sodium bicarbonate solution (25 ml) was added. Extraction with ethyl acetate, followed by solvent evaporation *in vacuo* gave 2-(4-amino-1*H*-1,2,3-triazol-1-yl)-*N*-(3-fluorophenyl)acetamide as a beige solid (1.0 g, 95 % yield) :

-52-

¹H-NMR (DMSO d₆) : 10.60 (s, 1H), 7.55 (m, 1H), 7.37 (m, 1H), 7.3 (m, 1H), 7.15 (s, 1H), 6.92 (m, 1H), 5.13 (s, 2H), 4.73 (s, 2H) :

MS (+ve ESI): 236.2 (M+H)⁺.

4-chloro-7-(3-chloropropoxy)-6-methoxyquinazoline, used as starting material, was obtained as follows:

- f) Palladium on carbon (3.3 g of a 10 % mixture) was added to a solution of 7-(benzyloxy)-6-methoxyquinazolin-4-(3H)-one (20 g, 71 mmol) (prepared according to *J. Med. Chem.* **1999**, 42, 5369-5389) suspended in dimethylformamide (530 ml). Ammonium formate (45 g, 710 mmol) was then added portion-wise over 1.25 hour. The reaction mixture was stirred for an additional 0.5 hour and the catalyst was removed by filtration. The solvent was removed *in vacuo* to yield 7-hydroxy-6-methoxyquinazolin-4-(3H)-one (8.65 g, 64 % yield) : ¹H-NMR (DMSO d₆) : 7.91 (s, 1H), 7.45 (s, 1H), 7.01 (s, 1H), 3.90 (s, 3H).
- g) A mixture of 7-hydroxy-6-methoxyquinazolin-4-(3H)-one (8.0 g, 41.6 mmol), pyridine (7.5 ml) and acetic anhydride (63 ml) was heated at 100°C for 4.5 hours and left to cool to ambient temperature for 18 hours. The reaction mixture was poured into ice/water (400 ml) and the resultant precipitate collected by filtration and dried *in vacuo*. Analysis revealed that hydrolysis of the acetate group on the 4 position of the quinazoline was incomplete. The mixture was therefore treated with water (150 ml) and pyridine (0.5 ml) at 90 °C for 15 minutes. The reaction was cooled and the solid was collected by filtration, washed with water and dried *in vacuo* to yield 7-(acetoxy)-6-methoxyquinazolin-4-(3H)-one (7.4 g, 76 % yield) : ¹H-NMR (DMSO d₆) : 8.05 (s, 1H), 7.65 (s, 1H), 7.45 (s, 1H), 3.90 (s, 3H), 2.31 (s, 3H).
- h) Dimethylformamide (0.5 ml) was added to a solution of 7-(acetoxy)-6-methoxyquinazolin-4-(3H)-one (2.0 g, 8.5 mmol) in thionyl chloride (32 ml) and the reaction mixture was heated at reflux for 1.5 hours. Upon cooling to ambient temperature, the thionyl chloride was removed *in vacuo* and azeotroped with toluene. The residue was diluted with dichloromethane (15 ml), a solution of 10 % ammonia in methanol (80 ml) added and the mixture heated at 80°C for 10 minutes. Upon cooling to ambient temperature, the solvent was evaporated to almost complete dryness, water was added and the pH adjusted to 7 with dilute hydrochloric acid. The resultant precipitate was collected by filtration and dried *in vacuo* at 30 °C for 18 hours to yield 4-chloro-7-hydroxy-6-methoxyquinazoline (1.65 g, 92 % yield) : ¹H-NMR (DMSO d₆) : 8.81 (s, 1H), 7.40 (s, 1H), 7.25 (s, 1H), 4.00 (s, 3H).
- i) Triphenylphosphine (2.6 g, 10.1 mmol) and 3-chloropropanol (0.69 ml, 8.2 mmol) were added to a suspension of 4-chloro-7-hydroxy-6-methoxyquinazoline (1.65 g, 7.8 mmol)

-53-

in dichloromethane (100 ml) under argon. The flask was placed in a water bath at 20 °C and di-*tert*-butyl azodicarboxylate (2.30 g, 10.1 mmol) added portion wise over a few minutes. The reaction mixture was stirred at ambient temperature for 2 hours before solvent evaporation *in vacuo*. Purification by flash chromatography on silica gel, eluting with ethyl acetate : petroleum ether (3:7) yielded 4-chloro-7-(3-chloropropoxy)-6-methoxyquinazoline (2.0 g, 91 % yield) :

¹H-NMR (DMSO d₆) : 8.90 (s, 1H), 7.55 (s, 1H), 7.45 (s, 1H), 4.42 (m, 2H), 4.05 (s, 3H), 3.80 (m, 2H), 2.31 (m, 2H).

10 **Example 2 - Preparation of compound 2 in table 1 - 2-(4-{[7-(3-chloropropoxy)quinazolin-4-yl]amino}-1H-1,2,3-triazol-1-yl)-N-(3-fluorophenyl)acetamide**

2-(4-amino-1H-1,2,3-triazol-1-yl)-N-(3-fluorophenyl)acetamide (446 mg, 1.9 mmol) was added to a solution of 4-chloro-7-(3-chloropropoxy)quinazoline

15 (488 mg, 1.9 mmol) in dimethylacetamide (15 ml). A solution of hydrochloric acid in dioxane (4.0 N, 475 µl, 1.9 mmol) was added to the reaction mixture and the resulting solution was heated at 90 °C for 3 hours. The mixture was cooled, diluted with isopropanol and the solid recovered by suction filtration. Washing the solid with ethyl acetate and diethyl ether, followed by prolonged drying *in vacuo*, gave compound 2 in table 1 (620 mg, 66 % yield) :

20 ¹H-NMR (DMSO d₆, TFA) : 9.10 (s, 1H), 8.92 (d, 1H), 8.72 (s, 1H), 7.61 (m, 1H), 7.54 (m, 1H), 7.39 (m, 3H), 6.93 (t, 1H), 5.51 (s, 2H), 4.37 (t, 2H), 3.86 (t, 2H), 2.31 (m, 2H) :

MS (+ve ESI): 456.1 (M+H)⁺

4-chloro-7-(3-chloropropoxy)quinazoline, used as the starting material was obtained as follows:

25 a) Formamidinium acetate (20.13 g, 193.4 mmol) was added to a solution of 2-amino-4-fluorobenzoic acid (15.0 g, 96.7 mmol) in 2-methoxyethanol (97 ml) and the mixture heated to reflux for 18 hours. The reaction was cooled, concentrated and the residue stirred in aqueous ammonium hydroxide (0.01 N, 250 ml) for 1 hour. The suspension was filtered, washed with water and dried over phosphorus pentoxide to yield 7-fluoroquinazolin-4-ol as an
30 off-white solid (10.35 g, 65 % yield) :

¹H-NMR (DMSO d₆) : 12.32 (br s, 1H), 8.19 (dd, 1H), 8.14 (s, 1H), 7.45 (m, 1H), 7.39 (m, 1H) :

¹⁹F-NMR (DMSO d₆): -105 (m) :

MS (-ve ESI): 163 (M-H)⁻,

MS (+ve ESI): 165 (M+H)⁺.

b) Sodium hydride (14.6 g, 365 mmol) was added at 0 °C to a solution of 1,3-propanediol (27.8 g, 365 mmol) in dimethylformamide (70 ml). 7-fluoroquinazolin-4-ol (10 g, 60.9 mmol) was added portion-wise and the reaction mixture heated at 60 °C, then at 100 °C for 3 hours. The reaction was cooled to 0 °C, quenched with water (280 ml) and adjusted to pH 5.9. The resulting suspension was filtered, washed with water then diethyl ether and dried over phosphorus pentoxide to yield 7-(3-hydroxypropoxy)quinazolin-4-ol as a white powder (12.4 g, 92 % yield) :

¹H-NMR (DMSO d₆) : 11.90 (br s, 1H), 8.04 (s, 1H), 8.00 (d, 1H), 7.10 (m, 2H), 4.17 (t, 2H), 3.58 (t, 2H), 1.92 (m, 2H) :

MS (+ve ESI): 221 (M+H)⁺.

c) Dimethylformamide (1 ml) was added to a mixture of 7-(3-hydroxypropoxy)quinazolin-4-ol (10.5 g, 47.7 mmol) and thionyl chloride (100 ml, 137 mmol) and the reaction mixture heated to 85 °C for 1 hour. The mixture was cooled to ambient temperature, diluted with toluene and evaporated to dryness. This was repeated until all thionyl chloride was removed. The residue was dissolved in dichloromethane and washed with a saturated sodium bicarbonate solution. The aqueous layer was extracted with dichloromethane and the combined organics were dried (magnesium sulphate) and concentrated to leave a yellow solid. Trituration with diethyl ether removed a less soluble impurity and the diethyl ether filtrate was concentrated to yield 4-chloro-7-(3-chloropropoxy)quinazoline as an off-white solid (8.5 g, 70 % yield) :

¹H-NMR (DMSO d₆) : 13.25 (br s, 1H), 8.34 (s, 1H), 8.06 (d, 1H), 7.17 (m, 2H), 4.21 (t, 2H), 3.83 (t, 2H), 2.23 (m, 2H).

MS (+ve ESI): 257, 259 (M+H)⁺.

Example 3 - Preparation of compound 3 in table 1 -(4-{[7-(3-chloropropoxy)quinazolin-4-yl]amino}-1H-1,2,3-triazol-1-yl)-N-(2,3-difluorophenyl)acetamide

An analogous reaction to that described in example 2, but starting with 2-(4-amino-1H-1,2,3-triazol-1-yl)-N-(2,3-difluorophenyl)acetamide (2.1 g, 8.3 mmol) yielded compound 3 in table 1 (3.9 g, 92 % yield):

¹H-NMR (DMSO d₆, TFA): 9.06 (s, 1H), 8.87 (d, 1H), 8.68 (s, 1H), 7.71 (m, 1H), 7.5 (d, 1H), 7.34 (s, 1H), 7.18 (m, 2H), 5.57 (s, 2H), 4.33 (t, 2H), 3.83 (t, 2H), 2.27 (m, 2H).

MS (+ve ESI): 474.15 (M+H)⁺.

2-(4-amino-1*H*-1,2,3-triazol-1-yl)-*N*-(2,3-difluorophenyl)acetamide used as starting material, was prepared as follows:

- a) An analogous reaction to that described in example 1d, but starting with 2,3 difluoroaniline (5 ml, 49 mmol) yielded *tert*-butyl (1-{2-[(2,3-difluorophenyl)amino]-2-oxoethyl}-1*H*-1,2,3-triazol-4-yl)carbamate (11.2 g, 78 % yield):

¹H-NMR (DMSO-d₆): 10.47 (s, 1H), 10.04 (brs, 1H), 7.96 (s, 1H), 7.7 (t, 1H), 7.21 (m, 2H), 5.37 (s, 2H), 1.46 (s, 9H).

MS (+ve ESI): 354.2 (M+H)⁺.

- b) An analogous reaction to that described in example 1e, but starting with *tert*-butyl (1-{2-[(2,3-difluorophenyl)amino]-2-oxoethyl}-1*H*-1,2,3-triazol-4-yl)carbamate (11.1 g, 31 mmol) yielded 2-(4-amino-1*H*-1,2,3-triazol-1-yl)-*N*-(2,3-difluorophenyl)acetamide (3 g, 39 % yield):

¹H-NMR (DMSO d₆): 10.41 (s, 1H), 7.7 (t, 1H), 7.21 (m, 2H), 7.14 (s, 1H), 5.22 (s, 1H), 4.73 (s, 2H).

MS (+ve ESI): 254.21 (M+H)⁺.

Example 4 - Preparation of compound 4 in table 2 - *N*-(3-fluorophenyl)-2-{4-[(7-{3-[(2-hydroxyethyl)(propyl)amino]propoxy}-6-methoxyquinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl}acetamide

2-(4-{[7-(3-chloropropoxy)-6-methoxyquinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)-*N*-(3-fluorophenyl)acetamide (137 mg, 0.23 mmol) was added to a solution of 2-(propylamino)ethanol (95 mg, 0.92 mmol) in dimethylacetamide (0.5 ml) in the presence of potassium iodide (76 mg, 0.46 mmol) and the reaction was heated under argon at 95 °C for 3 hours. The reaction was cooled, the solvent was evaporated *in vacuo* and the residue was purified by preparative LCMS. The fractions containing the desired compound were combined, evaporated *in vacuo* and the residue was dissolved in a mixture of dichloromethane (5 ml) and methanol (5 ml). Addition of a small volume of diethyl ether caused precipitation of a solid which was collected by suction filtration and dried *in vacuo* to give compound 4 in table 2 (75 mg, 55 % yield) :

¹H-NMR (DMSO d₆, TFA) : 9.07 (s, 1H), 8.72 (s, 1H), 8.39 (s, 1H), 7.60 (d, 1H), 7.40 (m, 2H), 7.34 (d, 1H), 6.95 (t, 1H), 5.50 (s, 2H), 4.38 (m, 1H), 4.32 (m, 2H), 4.03 (s, 3H), 3.78

-56-

(m, 1H), 3.53 (m, 1H), 3.37 (m, 2H), 3.28 (m, 1H), 3.18 (m, 2H), 2.29 (m, 2H), 1.72 (m, 2H), 0.95 (m, 3H) :

MS (+ve ESI): 553.3 (M+H)⁺.

5 **Example 5 - Preparation of compound 5 in table 2 - N-(3-fluorophenyl)-2-{4-[(7-{3-[(2S)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy}-6-methoxyquinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide**

An analogous reaction to that described in example 4, but starting with (2S)-pyrrolidin-2-ylmethanol (93 mg, 0.98 mmol), yielded compound 5 in table 2 (90 mg, 66 %
10 yield) :

¹H-NMR (DMSO d₆, TFA) : 9.07 (s, 1H), 8.73 (s, 1H), 8.39 (s, 1H), 7.60 (m, 1H), 7.40 (m, 2H), 7.34 (m, 1H), 6.94 (m, 1H), 5.51 (s, 1H), 4.33 (m, 2H), 4.03 (s, 3H), 3.77 (m, 1H), 3.59 (m, 4H), 3.25 (m, 2H), 2.31 (m, 2H), 2.13 (m, 1H), 2.04 (m, 1H), 1.90 (m, 1H), 1.79 (m, 1H) :
MS (+ve ESI) : 551.3 (M+H)⁺.

15

Example 6 - Preparation of compound 6 in table 2 - N-(3-fluorophenyl)-2-{4-[(7-{3-[(2S)-2-(hydroxyethyl)(propyl)amino]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

2-(4-[(7-(3-chloropropoxy)quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl)-N-(3-
20 fluorophenyl)acetamide (138 mg, 2.8 mmol) was added to a solution of 2-(propylamino)ethanol (115 mg, 11.2 mmol) in dimethylacetamide (0.5 ml) in the presence of potassium iodide (93 mg, 5.6 mmol) and the reaction heated under argon at 90 °C for 3 hours. The reaction was cooled, the solvent was evaporated *in vacuo* and the residue was purified by preparative LCMS. The fractions containing the desired compound were combined,
25 evaporated *in vacuo* and the residue was dissolved in a mixture of dichloromethane (5 ml) and methanol (5 ml). Addition of a small volume of diethyl ether caused precipitation of a solid which was collected by suction filtration and dried *in vacuo* to give compound 6 in table 2 (85 mg, 58 % yield) :

¹H-NMR (DMSO d₆, TFA) : 9.11 (s, 1H), 8.92 (d, 1H), 8.72 (s, 1H), 7.60 (d, 1H), 7.53 (m, 1H), 7.40 (m, 1H), 7.34 (m, 2H), 6.94 (t, 1H), 5.51 (s, 2H), 4.33 (t, 2H), 3.79 (t, 2H), 3.35 (m, 2H), 3.27 (m, 2H), 3.15 (m, 2H), 2.26 (m, 2H), 1.73 (m, 2H), 0.95 (m, 3H) :
30 MS (+ve ESI): 523.0 (M+H)⁺.

Example 7 - Preparation of compound 7 in table 2 - N-(3-fluorophenyl)-2-{4-[(7-{3-[(2S)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

An analogous reaction to that described in example 6, but starting with (2S)-
5 pyrrolidin-2-ylmethanol (105 mg, 1.12 mmol), yielded compound 7 in table 2 (60 mg, 41 % yield) :

¹H-NMR (DMSO d₆, TFA) : 9.10 (s, 1H), 8.91 (d, 1H), 8.71 (s, 1H), 7.60 (d, 1H), 7.53 (m, 1H), 7.40 (m, 1H), 7.34 (m, 2H), 6.94 (t, 1H), 5.51 (s, 2H), 4.33 (t, 2H), 3.78 (m, 1H), 3.63 (m, 4H), 3.27 (m, 1H), 3.19 (m, 1H), 2.27 (m, 2H), 2.14 (m, 1H), 2.04 (m, 1H), 1.91 (m, 1H),
10 1.80 (m, 1H) :

MS (+ve ESI): 521.0 (M+H)⁺.

Example 8 - Preparation of compound 8 in table 2 - N-(3-fluorophenyl)-2-(4-{[7-(3-morpholin-4-yl)propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl)acetamide

15 An analogous reaction to that described in example 6 but starting with morpholine (105 mg, 1.2 mmol) yielded compound 8 in table 2 (55 mg, 36% yield):

¹H-NMR (DMSO d₆, TFA): 9.12 (s, 1H); 8.94 (d, 1H); 8.73 (s, 1H); 7.62 (ddd, 1H); 7.53 (dd, 1H); 7.40 (dd, 1H); 7.39-7.32 (m, 2H); 6.93 (ddd, 1H); 5.52 (s, 2H); 4.35 (t, 2H); 4.05 (dd, 2H); 3.73 (dd, 2H); 3.56 (d, 2H); 3.44-3.34 (m, 2H); 3.24-3.13 (m, 2H); 2.34-2.25 (m, 2H).
20 MS (+ve ESI): 507.2 (M+H)⁺.

Example 9 - Preparation of compound 9 in table 2 - N-(3-fluorophenyl)-2-(4-{[7-(3-piperidin-1-yl)propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl)acetamide

An analogous reaction to that described in example 6 but starting with piperidine (102
25 mg, 1.2 mmol) yielded compound 9 in table 2 (26 mg, 17% yield):

¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.93 (d, 1H); 8.73 (s, 1H); 7.61 (ddd, 1H); 7.52 (dd, 1H); 7.40 (dd, 1H); 7.37-7.32 (m, 2H); 6.93 (ddd, 1H); 5.51 (s, 2H); 4.33 (t, 2H); 3.54 (d, 2H); 3.33-3.24 (m, 2H); 3.01-2.90 (m, 2H); 2.32-2.21 (m, 2H); 1.92-1.82 (m, 2H); 1.79-1.63 (m, 3H); 1.49-1.37 (m, 1H).
30 MS (+ve ESI): 504.6 (M+H)⁺.

Example 10 - Preparation of compound 10 in table 2 - N-(3-fluorophenyl)-2-(4-{[7-(3-pyrrolidin-1-yl)propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl)acetamide

-58-

An analogous reaction to that described in example 6 but starting with pyrrolidine (85 mg, 1.2 mmol) yielded compound 10 in table 2 (43 mg, 29% yield):

¹H-NMR (DMSO d₆, TFA): 9.12 (s, 1H); 8.93 (d, 1H); 8.73 (s, 1H); 7.61 (ddd, 1H); 7.53 (dd, 1H); 7.40 (dd, 1H); 7.37-7.32 (m, 2H); 6.93 (ddd, 1H); 5.51 (s, 2H); 4.33 (t, 2H); 3.70-3.61 (m, 2H); 3.43-3.35 (m, 2H); 3.15-3.04 (m, 2H); 2.31-2.20 (m, 2H); 2.13-2.02 (m, 2H); 1.96-1.87 (m, 2H).

MS (+ve ESI): 491.2 (M+H)⁺.

Example 11 - Preparation of compound 11 in table 2 - N-(3-fluorophenyl)-2-{4-[(7-{3-[(2-hydroxy-1,1-dimethylethyl)amino]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

An analogous reaction to that described in example 6 but starting with 2-amino-2-methylpropan-1-ol (107 mg, 1.2 mmol) yielded compound 11 in table 2 (80 mg, 52 % yield):

¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.61 (ddd, 1H); 7.54 (dd, 1H); 7.40 (dd, 1H); 7.37-7.32 (m, 2H); 6.94 (ddd, 1H); 5.51 (s, 2H); 4.35 (s, 2H); 3.47 (s, 2H); 3.15-3.07 (m, 2H); 2.27-2.17 (m, 2H); 1.26 (s, 6H).

MS (+ve ESI): 508.6 (M+H)⁺.

Example 12 - Preparation of compound 12 in table 2 - 2-[4-[(7-{3-(cyclopropylamino)propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl]-N-(3-fluorophenyl)acetamide

An analogous reaction to that described in example 6 but starting with cyclopropylamine (69 mg, 1.2 mmol) yielded compound 12 in table 2 (25 mg, 17 % yield):

¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.61 (ddd, 1H); 7.54 (dd, 1H); 7.40 (dd, 1H); 7.37-7.31 (m, 2H); 6.94 (ddd, 1H); 5.51 (s, 2H); 4.34 (t, 2H); 3.31-3.21 (m, 2H); 2.85-2.76(m, 1H); 2.26-2.16 (m, 2H); 0.91-0.77 (m, 4H).

MS (+ve ESI): 477.2 (M+H)⁺.

Example 13 - Preparation of compound 13 in table 2 - 2-{4-[(7-{3-[(2-dimethylamino)ethyl](methyl)amino]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}-N-(3-fluorophenyl)acetamide

-59-

An analogous reaction to that described in example 6 but starting with N,N,N'-trimethylethane-1,2-diamine (123 mg, 1.2 mmol) yielded compound 13 in table 2 (87 mg, 56 % yield):

¹H-NMR (DMSO d₆, TFA): 9.12 (s, 1H); 8.94 (d, 1H); 8.73 (s, 1H); 7.62 (ddd, 1H); 7.53 (dd, 1H); 7.44-7.33 (m, 3H); 6.93 (ddd, 1H); 5.52 (2H); 4.35 (t, 2H); 3.71-3.50 (m, 4H); 3.48-3.36 (m, 2H); 2.95 (s, 3H); 2.92 (s, 6H); 2.36-2.24 (m, 2H).

MS (+ve ESI): 522.3 (M+H)⁺.

Example 14 - Preparation of compound 14 in table 2 - N-(3-fluorophenyl)-2-[4-({7-[3-(4-methylpiperazin-1-yl)propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl]acetamide

An analogous reaction to that described in example 6 but starting with 1-methylpiperazine (120 mg, 1.2 mmol) yielded compound 14 in table 2 (83 mg, 53 % yield):

¹H-NMR (DMSO d₆, TFA): 9.12 (s, 1H); 8.93 (d, 1H); 8.73 (s, 1H); 7.61 (ddd, 1H); 7.53 (dd, 1H); 7.43-7.32 (m, 3H); 6.93 (ddd, 1H); 5.52 (s, 2H); 4.35 (t, 2H); 3.52-3.42 (m, 2H); 4.08-3.11 (m, 8H); 2.97 (s, 3H); 2.33-2.23 (m, 2H).

MS (+ve ESI): 520.3 (M+H)⁺.

Example 15 - Preparation of compound 15 in table 2 - N-(3-fluorophenyl)-2-[4-[(7-{3-[(2R)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl]acetamide

An analogous reaction to that described in example 6 but starting with (2R)-pyrrolidin-2-ylmethanol (121 mg, 1.2 mmol) yielded compound 15 in table 2 (120 mg, 77% yield):

¹H-NMR (DMSO d₆, TFA): 9.12 (s, 1H); 8.94 (d, 1H); 8.73 (s, 1H); 7.61 (ddd, 1H); 7.53 (dd, 1H); 7.40 (dd, 1H); 7.38-7.33 (m, 2H); 6.93 (ddd, 1H); 5.52 (s, 2H); 4.34 (t, 2H); 3.84-3.77 (m, 1H); 3.70-3.56 (m, 4H); 3.33-3.25 (m, 1H); 3.24-3.15 (1H); 2.33-2.24 (m, 2H); 2.20-2.11 (m, 1H); 2.09-2.00 (m, 1H); 1.96-1.87 (m, 1H); 1.85-1.75 (m, 1H).

MS (+ve ESI): 521.2 (M+H)⁺.

Example 16 - Preparation of compound 16 in table 2 - N-(3-fluorophenyl)-2-[4-({7-[3-(4-hydroxypiperidin-1-yl)propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl]acetamide

An analogous reaction to that described in example 6 but starting with piperidin-4-ol (121 mg, 1.2 mmol) yielded compound 16 in table 2 (130 mg, 83 % yield):

-60-

¹H-NMR (DMSO *d*₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.73 (s, 1H); 7.61 (ddd, 1H); 7.53 (ddd, 1H); 7.40 (dd, 1H); 7.37-7.32 (m, 2H); 6.93 (ddd, 1H); 5.51 (s, 2H); 4.37-4.29 (m, 2H); 4.02-3.96 (m, 0.5H); 3.73-3.64 (m, 0.5H); 3.60-3.51 (m, 1H); 3.44-3.16 (m, 4H); 3.09-2.98 (m, 1H); 2.31-2.21 (m, 2H); 2.07-1.99 (m, 1H); 1.94-1.77 (m, 2H); 1.55-1.67 (m, 1H).

5 MS (+ve ESI): 521.2 (M+H)⁺.

Example 17 - Preparation of compound 17 in table 2 - 2-{4-[(7-{3-[ethyl(2-hydroxyethyl)amino]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl]-N-(3-fluorophenyl)acetamide

10 An analogous reaction to that described in example 6 but starting with 2-(ethylamino)ethanol (107 mg, 1.2 mmol) yielded compound 17 in table 2 (112 mg, 73 % yield):

¹H-NMR (DMSO *d*₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.61 (ddd, 1H); 7.54 (dd, 1H); 7.40 (dd, 1H); 7.37-7.32 (m, 2H); 6.93 (ddd, 1H); 5.51 (s, 2H); 4.38-4.29 (m, 2H); 3.83-
15 3.74 (m, 2H); 3.43-3.20 (m, 6H); 2.33-2.19 (m, 2H); 1.27 (t, 3H).

MS (+ve ESI): 509.2 (M+H)⁺.

Example 18 - Preparation of compound 18 in table 2 - N-(3-fluorophenyl)-2-{4-[(7-{3-[4-(2-hydroxyethyl)piperazin-1-yl]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

20 An analogous reaction to that described in example 6 but starting with 2-piperazin-1-yl-ethanol (156 mg, 1.2 mmol) yielded compound 18 in table 2 (132 mg, 80 % yield):

¹H-NMR (DMSO *d*₆, TFA): 9.12 (s, 1H); 8.93 (d, 1H); 8.73 (s, 1H); 7.61 (ddd, 1H); 7.53 (dd, 1H); 7.47-7.42 (m, 3H); 6.93 (ddd, 1H); 5.52 (s, 2H); 3.40-3.31 (m, 2H); 3.84-3.77 (m, 2H);
25 3.51-3.43 (m, 2H); 3.42-3.34 (m, 2H); 4.07-3.25 (m, 8H); 2.36-2.24 (m, 2H).

MS (+ve ESI): 550.3 (M+H)⁺.

Example 19 - Preparation of compound 19 in table 2 - N-(3-fluorophenyl)-2-(4-[(7-(3-piperazin-1-ylpropoxy)quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl)acetamide

30 An analogous reaction to that described in example 6 but starting with *tert*-butyl piperazine-1-carboxylate (224 mg, 1.2 mmol) yielded compound 19 in table 2 (88 mg, 58 % yield) after treatment with hydrochloric acid in diethyl ether:

-61-

¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.61 (ddd, 1H); 7.53 (dd, 1H); 7.44-7.32 (m, 3H); 6.94 (ddd, 1H); 5.51 (s, 2H); 4.39-4.30 (m, 2H); 3.49-3.41 (m, 2H); 4.10-2.90 (m, 8H); 2.35-2.23 (m, 2H).

MS (+ve ESI): 506.2 (M+H)⁺.

5

Example 20 - Preparation of compound 20 in table 2 - N-(3-fluorophenyl)-2-{4-[(7-{3-[4-(2-hydroxyethyl)piperidin-1-yl]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

An analogous reaction to that described in example 6 but starting with 2-piperidin-4-yl-ethanol (155 mg, 1.2 mmol) yielded compound 20 in table 2 (111 mg, 67 % yield):

¹H-NMR (DMSO d₆, TFA): 9.12 (s, 1H); 8.94 (d, 1H); 8.74 (s, 1H); 7.62 (ddd, 1H); 7.52 (dd, 1H); 7.43-7.33 (m, 3H); 6.92 (ddd, 1H); 5.52 (s, 2H); 4.37-4.30 (m, 2H); 3.62-3.55 (m, 2H); 3.54-3.47 (m, 2H); 3.34-3.26 (m, 2H); 3.05-2.93 (m, 2H); 2.34-2.22 (m, 2H); 1.99-1.89 (m, 2H); 1.79-1.66 (m, 1H); 1.47-1.37 (m, 4H).

MS (+ve ESI): 549.3 (M+H)⁺.

Example 21 - Preparation of compound 21 in table 2 - N-(3-fluorophenyl)-2-{4-[(7-{3-[4-(hydroxymethyl)piperidin-1-yl]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

An analogous reaction to that described in example 6 but starting with piperidin-4-ylmethanol (138 mg, 1.2 mmol) yielded compound 21 in table 2 (74 mg, 46 % yield):

¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.93 (d, 1H); 8.72 (s, 1H); 7.61 (ddd, 1H); 7.52 (dd, 1H); 7.40 (dd, 1H); 7.37-7.32 (m, 2H); 6.93 (ddd, 1H); 5.51 (s, 2H); 4.33 (t, 2H); 3.66-3.55 (m, 2H); 3.39-3.22 (m, 4H); 3.05-2.91 (m, 2H); 2.33-2.20 (m, 2H); 1.95-1.85 (m, 2H); 1.76-1.62 (m, 1H); 1.51-1.37 (m, 2H).

MS (+ve ESI): 535.3 (M+H)⁺.

Example 22 - Preparation of compound 22 in table 2 - N-(3-fluorophenyl)-2-{4-[(7-{3-[(2-hydroxyethyl)(isopropyl)amino]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

An analogous reaction to that described in example 6 but starting with 2-(isopropylamino)ethanol (124 mg, 1.2 mmol) yielded compound 22 in table 2 (92 mg, 59 % yield):

-62-

¹H-NMR (DMSO *d*₆, TFA): 9.12 (s, 1H); 8.94 (d, 1H); 8.73 (s, 1H); 7.62 (ddd, 1H); 7.51 (dd, 1H); 7.40 (dd, 1H); 7.38-7.33 (m, 2H); 6.93 (ddd, 1H); 5.52 (s, 2H); 4.40-4.30 (m, 2H); 3.85-3.70 (m, 3H); 3.41-3.28 (m, 3H); 3.23-3.13 (m, 1H); 1.32 (d, 3H); 1.31 (d, 3H).

MS (+ve ESI): 523.3 (M+H)⁺.

5

Example 23 - Preparation of compound 23 in table 2 - 2-{4-[(7-{3-[cyclopropyl(2-hydroxyethyl)amino]propoxy}quinazolin-4-yl)amino]-1*H*-1,2,3-triazol-1-yl]}-*N*-(3-fluorophenyl)acetamide

An analogous reaction to that described in example 6 but starting with 2-

10 (cyclopropylamino)ethanol (121 mg, 1.2 mmol) yielded compound 23 in table 2 (73 mg, 47 % yield):

¹H-NMR (DMSO *d*₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.60 (ddd, 1H); 7.54 (dd, 1H); 7.40 (dd, 1H); 7.37-7.30 (m, 2H); 6.94 (ddd, 1H); 5.51 (s, 2H); 4.38-4.30 (m, 2H); 3.93-3.75 (m, 2H); 3.57-3.35 (m, 4H); 2.98-2.89 (m, 1H); 2.40-2.27 (m, 2H); 1.12-0.84 (m, 4H).

15 MS (+ve ESI): 521.3 (M+H)⁺.

Example 24 - Preparation of compound 24 in table 2 - *N*-(2,3-difluorophenyl)-2-(4-{[7-(3-morpholin-4-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide

An analogous reaction to that described in example 6 but starting with morpholine

20 (105 mg, 1.2 mmol) and 2-(4-{[7-(3-chloropropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)-*N*-(2,3-difluorophenyl)acetamide (153 mg, 0.3 mmol) yielded compound 24 in table 2 (115 mg, 73 % yield):

¹H-NMR (DMSO *d*₆, TFA): 9.11 (s, 1H); 8.93 (d, 1H); 8.72 (s, 1H); 7.80-7.72 (m, 1H); 7.52 (dd, 1H); 7.35 (d, 1H); 7.26-7.16 (m, 2H); 5.60 (s, 2H); 4.34 (t, 2H); 4.08-4.01 (m, 2H); 3.76-3.66 (m, 2H); 3.59-3.51 (m, 2H); 3.41-3.34 (m, 2H); 3.21-3.11 (m, 2H); 2.33-2.23 (m, 2H).
25 MS (+ve ESI): 525.2 (M+H)⁺.

Example 25 - Preparation of compound 25 in table 2 - *N*-(2,3-difluorophenyl)-2-(4-{[7-(3-piperidin-1-ylpropoxy)quinazolin-4-yl]amino}-1*H*-1,2,3-triazol-1-yl)acetamide

30 An analogous reaction to that described in example 24 but starting with piperidine (102 mg, 1.2 mmol) yielded compound 25 in table 2 (101 mg, 65 % yield):

¹H-NMR (DMSO *d*₆, TFA): 9.11 (s, 1H); 8.91 (d, 1H); 8.72 (s, 1H); 7.79-7.71 (m, 1H); 7.52 (dd, 1H); 7.35 (d, 1H); 7.27-7.16 (m, 2H); 5.60 (s, 2H); 4.36-4.28 (m, 2H); 3.57-3.48 (m, 2H);

-63-

3.31-3.24 (m, 2H); 3.00-2.90 (m, 2H); 2.30-2.21 (m, 2H); 1.91-1.82 (m, 2H); 1.78-1.62 (m, 3H); 1.49-1.37 (m, 1H).

MS (+ve ESI): 523.2 (M+H)⁺.

5 **Example 26 - Preparation of compound 26 in table 2 - N-(2,3-difluorophenyl)-2-(4-{[7-(3-pyrrolidin-1-ylpropoxy)quinazolin-4-yl]amino}-1H-1,2,3-triazol-1-yl)acetamide**

An analogous reaction to that described in example 24 but starting with pyrrolidine (85 mg, 1.2 mmol) yielded compound 26 in table 2 (50 mg, 33 % yield):

1H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.79-7.71 (m, 1H); 7.53 (dd, 1H); 7.34 (d, 1H); 7.27-7.16 (m, 2H); 5.60 (s, 2H); 4.33 (t, 2H); 3.69-3.59 (m, 2H); 3.42-3.33 (m, 2H); 3.14-3.03 (m, 2H); 2.29-2.18 (m, 2H); 2.13-2.00 (m, 2H); 1.98-1.85 (m, 2H).
MS (+ve ESI): 509.2 (M+H)⁺.

15 **Example 27 - Preparation of compound 27 in table 2 - N-(2,3-difluorophenyl)-2-{4-[(7-{3-[(2-hydroxy-1,1-dimethylethyl)amino]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide**

An analogous reaction to that described in example 24 but starting with 2-amino-2-methylpropan-1-ol (107 mg, 1.2 mmol) yielded compound 27 in table 2 (69 mg, 44 % yield):

1H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.79-7.72 (m, 1H); 7.54 (dd, 1H); 7.34 (d, 1H); 7.25-7.16 (m, 2H); 5.61 (s, 2H); 4.35 (t, 2H); 3.48 (s, 2H); 3.16-3.06 (m, 2H); 2.26-2.16 (m, 2H); 1.26 (s, 6H).

MS (+ve ESI): 527.2 (M+H)⁺.

25 **Example 28 - Preparation of compound 28 in table 2 - 2-[4-{[7-{3-(cyclopropylamino)propoxy}quinazolin-4-yl]amino}-1H-1,2,3-triazol-1-yl]-N-(2,3-difluorophenyl)acetamide**

An analogous reaction to that described in example 24 but starting with cyclopropylamine (69 mg, 1.2 mmol) yielded compound 28 in table 2 (62 mg, 42 % yield):

1H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.91 (d, 1H); 8.72 (s, 1H); 7.79-7.70 (m, 1H); 7.57 (dd, 1H); 7.34 (d, 1H); 7.25-7.15 (m, 2H); 5.60 (d, 2H); 4.38-4.29 (m, 2H); 3.30-3.22 (m, 2H); 2.84-2.76 (m, 1H); 2.25-2.16 (m, 2H); 0.91-0.77 (m, 4H)..

MS (+ve ESI): 495.2 (M+H)⁺.

-64-

Example 29 - Preparation of compound 29 in table 2 - N-(2,3-difluorophenyl)-2-{4-[(7-{3-[[2-(dimethylamino)ethyl](methyl)amino]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

An analogous reaction to that described in example 24 but starting with N,N,N'-trimethylethane-1,2-diamine (123 mg, 1.2 mmol) yielded compound 29 in table 2 (89 mg, 55 % yield):

¹H-NMR (DMSO-d₆, TFA): 9.11 (s, 1H); 8.93 (d, 1H); 8.72 (s, 1H); 7.79-7.71 (m, 1H); 7.53 (dd, 1H); 7.37 (bs, 1H); 7.27-7.15 (m, 2H); 5.60 (s, 2H); 4.39-4.29 (m, 2H); 3.67-3.49 (m, 4H); 3.48-3.34 (m, 2H); 2.94 (s, 3H); 2.90 (s, 6H); 2.35-2.19 (m, 2H).

MS (+ve ESI): 540.3 (M+H)⁺.

Example 30 - Preparation of compound 30 in table 2 - N-(2,3-difluorophenyl)-2-[4-({7-[3-(4-methylpiperazin-1-yl)propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl]acetamide

An analogous reaction to that described in example 24 but starting with 1-methylpiperazine (120 mg, 1.2 mmol) yielded compound 30 in table 2 (64 mg, 39 % yield):

¹H-NMR (DMSO-d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.80-7.70 (m, 1H); 7.53 (dd, 1H); 7.36 (d, 1H); 7.27-7.16 (m, 2H); 5.60 (s, 2H); 4.41-4.29 (m, 2H); 4.17-3.10 (m, 8H); 3.50-3.40 (m, 2H); 2.95 (s, 3H); 2.33-2.23 (m, 2H).

MS (+ve ESI): 538.3 (M+H)⁺.

Example 31 - Preparation of compound 31 in table 2 - N-(2,3-difluorophenyl)-2-{4-[(7-{3-[(2R)-2-(hydroxymethyl)pyrrolidin-1-yl]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

An analogous reaction to that described in example 24 but starting with (2R)-pyrrolidin-2-ylmethanol (121 mg, 1.2 mmol) yielded compound 31 in table 2 (91 mg, 56 % yield):

¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.80-7.71 (m, 1H); 7.53 (dd, 1H); 7.35 (d, 1H); 7.27-7.16 (m, 2H); 5.60 (s, 2H); 4.40-4.29 (m, 2H); 3.83-3.75 (m, 1H); 3.69-3.54 (m, 4H); 3.34-3.24 (m, 1H); 3.23-3.14 (m, 1H); 2.34-2.23 (m, 2H); 2.20-2.09 (m, 1H); 2.09-1.98 (m, 1H); 1.96-1.85 (m, 1H); 1.84-1.74 (m, 1H).

MS (+ve ESI): 539.2 (M+H)⁺.

Example 32 - Preparation of compound 32 in table 2 - N-(2,3-difluorophenyl)-2-[4-({7-[3-(4-hydroxypiperidin-1-yl)propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl]acetamide

- An analogous reaction to that described in example 24 but starting with piperidin-4-ol (121 mg, 1.2 mmol) yielded compound 32 in table 2 (72 mg, 45 % yield):
- ¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.81-7.71 (m, 1H); 7.52 (ddd, 1H); 7.34 (bs, 1H); 7.27-7.16 (m, 2H); 5.61 (s, 2H); 4.39-4.28 (m, 2H); 4.00-3.96 (m, 0.5H); 3.74-3.65 (m, 0.5H); 3.60-3.52 (m, 1H); 3.44-3.15 (m, 4H); 3.09-2.98 (m, 1H); 2.32-2.20 (m, 2H); 2.07-1.98 (m, 1H); 1.94-1.77 (m, 2H); 1.67-1.54 (m, 1H).
- MS (+ve ESI): 539.2 (M+H)⁺.

Example 33 - Preparation of compound 33 in table 2 - N-(2,3-difluorophenyl)-2-[4-({7-[3-[ethyl(2-hydroxyethyl)amino]propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl]acetamide

- An analogous reaction to that described in example 24 but starting with 2-(ethylamino)ethanol (107 mg, 1.2 mmol) yielded compound 33 in table 2 (89 mg, 56 % yield):
- ¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.79-7.72 (m, 1H); 7.54 (dd, 1H); 7.34 (dd, 1H); 7.26-7.16 (m, 2H); 5.60 (s, 2H); 4.34 (t, 2H); 3.81-3.75 (m, 2H); 3.43-3.21 (m, 6H); 2.31-2.19 (m, 2H); 1.27 (t, 3H).
- MS (+ve ESI): 527.2 (M+H)⁺.

Example 34 - Preparation of compound 34 in table 2 - N-(2,3-difluorophenyl)-2-[4-({7-[3-[4-(2-hydroxyethyl)piperazin-1-yl]propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl]acetamide

- An analogous reaction to that described in example 24 but starting with 2-piperazin-1-ylethanol (156 mg, 1.2 mmol) yielded compound 34 in table 2 (89 mg, 52 % yield):
- ¹H-NMR (DMSO d₆, TFA): 9.12 (s, 1H); 8.93 (d, 1H); 8.73 (s, 1H); 7.80-7.72 (m, 1H); 7.53 (dd, 1H); 7.37 (dd, 1H); 7.26-7.16 (m, 2H); 5.60 (s, 2H); 4.38-4.31 (m, 2H); 4.10-3.10 (m, 8H); 3.83-3.76 (m, 2H); 3.50 (3.42 (m, 2H); 3.41-3.34 (m, 2H); 2.34-2.24 (m, 2H).
- MS (+ve ESI): 568.3 (M+H)⁺.

Example 35 - Preparation of compound 35 in table 2 - N-(2,3-difluorophenyl)-2-(4-({7-[3-(piperazin-1-yl)propoxy]quinazolin-4-yl}amino)-1H-1,2,3-triazol-1-yl)acetamide

An analogous reaction to that described in example 24 but starting with *tert*-butyl piperazine-1-carboxylate (224 mg, 1.2 mmol) yielded compound 35 in table 2 (96 mg, 61 % yield) after treatment with hydrochloric acid in diethyl ether:

¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.78-7.71 (m, 1H); 7.53 (dd, 1H); 7.36 (d, 1H); 7.27-7.16 (m, 2H); 5.60 (s, 2H); 4.39-4.29 (m, 2H); 4.20-3.00 (m, 8H); 3.49-3.40 (2H); 2.33-2.22 (m, 2H).

MS (+ve ESI): 524.3 (M+H)⁺.

10 **Example 36 - Preparation of compound 36 in table 2 - N-(2,3-difluorophenyl)-2-{4-[(7-{3-[4-(2-hydroxyethyl)piperidin-1-yl]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide**

An analogous reaction to that described in example 24 but starting with 2-piperidin-4-ylethanol (155 mg, 1.2 mmol) yielded compound 36 in table 2 (114 mg, 67 % yield):

¹H-NMR (DMSO d₆, TFA): 9.10 (s, 1H); 8.91 (d, 1H); 8.72 (s, 1H); 7.79-7.71 (m, 1H); 7.52 (dd, 1H); 7.34 (d, 1H); 7.28-7.16 (m, 2H); 5.60 (s, 2H); 4.32 (t, 2H); 3.60-3.52 (m, 2H); 3.48 (t, 2H); 3.32-3.22 (m, 2H); 3.01-2.93 (m, 2H); 2.30-2.21 (m, 2H); 1.95-1.86 (m, 2H); 1.76-1.62 (m, 1H); 1.45-1.32 (m, 4H).

MS (+ve ESI): 567.3 (M+H)⁺.

20 **Example 37 - Preparation of compound 37 in table 2 - N-(2,3-difluorophenyl)-2-{4-[(7-{3-[4-(hydroxymethyl)piperidin-1-yl]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide**

An analogous reaction to that described in example 24 but starting with piperidin-4-ylmethanol (138 mg, 1.2 mmol) yielded compound 37 in table 2 (73 mg, 44 % yield):

25 ¹H-NMR (DMSO d₆, TFA): 9.10 (s, 1H); 8.91 (d, 1H); 8.71 (s, 1H); 7.78-7.70 (m, 1H); 7.52 (dd, 1H); 7.34 (dd, 1H); 7.28-7.16 (m, 2H); 5.60 (s, 2H); 4.34-4.28 (m, 2H); 3.65-3.54 (m, 2H); 3.36-3.22 (m, 4H); 3.02-2.92 (m, 2H); 2.31-2.20 (m, 2H); 1.94-1.84 (m, 2H); 1.74-1.59 (m, 1H); 1.48-1.36 (m, 2H).

MS (+ve ESI): 553.3 (M+H)⁺.

30

Example 38 - Preparation of compound 38 in table 2 - N-(2,3-difluorophenyl)-2-{4-[(7-{3-[(2-hydroxyethyl)(isopropyl)amino]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}acetamide

-67-

An analogous reaction to that described in example 24 but starting with 2-(isopropylamino)ethanol (124 mg, 1.2 mmol) yielded compound 38 in table 2 (70 mg, 43 % yield):

¹H-NMR (DMSO d₆, TFA): 9.10 (s, 1H); 8.91 (d, 1H); 8.71 (s, 1H); 7.78-7.70 (m, 1H); 7.53 (dd, 1H); 7.34 (dd, 1H); 7.28-7.16 (m, 2H); 5.59 (s, 2H); 4.37-4.29 (m, 2H); 3.83-3.68 (m, 3H); 3.37-3.26 (m, 3H); 3.20-3.10 (m, 1H); 2.34-2.22 (m, 2H); 1.30 (d, 3H); 1.29 (d, 3H). MS (+ve ESI): 541.3 (M+H)⁺.

Example 39 - Preparation of compound 39 in table 2 - 2-{4-[(7-{3-[cyclopropyl(2-hydroxyethyl)amino]propoxy}quinazolin-4-yl)amino]-1H-1,2,3-triazol-1-yl}-N-(2,3-difluorophenyl)acetamide

An analogous reaction to that described in example 24 but starting with 2-(cyclopropylamino)ethanol (121 mg, 1.2 mmol) yielded compound 39 in table 2 (60 mg, 37 % yield):

¹H-NMR (DMSO d₆, TFA): 9.11 (s, 1H); 8.92 (d, 1H); 8.72 (s, 1H); 7.79-7.71 (m, 1H); 7.54 (dd, 1H); 7.36 (dd, 1H); 7.24-7.17 (m, 2H); 5.60 (s, 2H); 4.39-4.32 (m, 2H); 3.94-3.79 (m, 2H); 3.55-3.38 (m, 4H); 2.98-2.91 (m, 1H); 2.38-2.28 (m, 2H); 1.11-0.86 (m, 4H). MS (+ve ESI): 539.2 (M+H)⁺.